

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 0 481 214 B1

(12) EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention  
of the grant of the patent:  
24.06.1998 Bulletin 1998/26

(51) Int. Cl.<sup>6</sup>: C07F 9/6561, C07F 9/6512,  
C07F 9/6574, C07F 9/6584,  
A61K 31/675

(21) Application number: 91115312.0

(22) Date of filing: 10.09.1991

(54) Prodrugs of phosphonates

Wirkstoffvorläufer von Phosphonaten

Pro-médicaments dérivés de phosphonates

(84) Designated Contracting States:  
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(30) Priority: 14.09.1990 US 583906

(43) Date of publication of application:  
22.04.1992 Bulletin 1992/17

(73) Proprietors:  
• INSTITUTE OF ORGANIC CHEMISTRY AND  
BIOCHEMISTRY  
OF THE ACADEMY OF SCIENCES OF THE  
CZECH REPUBLIC  
166 10 Praha 6 (CZ)  
• Rega Stichting Vzw.  
B- 3000 Leuven (BE)

(72) Inventors:  
• Starrett, John Edward, Jr.  
Middletown, CT 06457 (US)  
• Mansuri, Muzammil M.  
Cheshire, CT 06410 (US)  
• Martin, John C.  
Cheshire, CT 06410 (US)  
• Tortolani, David R.  
Meriden, CT 06450 (US)

• Bronson, Joanne J.  
Madison, CT 06443 (US)

(74) Representative:  
Kinzebach, Werner, Dr. et al  
Patentanwälte  
Reitstötter, Kinzebach und Partner  
Postfach 86 06 49  
81633 München (DE)

(56) References cited:  
EP-A- 0 205 826 EP-A- 0 269 947  
EP-A- 0 270 885

• CHEMICAL ABSTRACTS vol. 112, no. 19, 7 May  
1990, abstract no. 179685h, Columbus, Ohio,  
US; A. HOLY et al.: 'Synthesis of N-(2-  
Phosphonylmethoxyethyl) Derivatives of  
Heterocyclic Bases'  
• CHEMICAL ABSTRACTS vol. 108, no. 21, 23 May  
1988, abstract no. 179636k, Columbus, Ohio, US;  
E. DE CLERCQ et al.: 'Antiviral Activity of  
Phosphonylmetoxyalkyl Derivatives of Purines  
and Pyrimidines'

EP 0 481 214 B1

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

## Description

The present invention relates to orally active prodrugs of phosphonate nucleotide analogs, their pharmaceutically acceptable acid addition salts, a process for their production, and to their use. The prodrugs of the present invention exhibit antitumor activity and a broad spectrum of antiviral activity.

Infectious viral diseases are recognized as an important medical problem. Progress against infectious viral diseases requires the development of drugs with selective antiviral activity while remaining benign to normal cell lines. Among the antiviral agents currently under study, which seem to possess selectivity, are phosphonate nucleotide analogs. In general, these compounds are structural analogs of the monophosphates nucleoside analogs.

A number of phosphonate nucleoside analogs have been described in the literature. These nucleoside analogs have been described as potent and selective antiviral agents with activity against a broad spectrum of DNA and RNA viruses.

For example, 9-(3-hydroxy-2-phosphonyl-methoxypropyl) (HPMP) and (2-phosphonyl-methoxy)ethyl (PME) analogs of purine (adenine (A), guanine (G), 2,6-diaminopurine (DAP), 2-monoaminopurine (MAP), hypoxanthine (Hx) and pyrimidine (cytosine (C), uracil (U), thymine (T)) were evaluated for antiviral properties. (S)-HPMPA, (S)-cyclic HPMPA, (S)-HPMPC, (S)-HPMPG, (S)-HPMPDAP, PMEDAP, PMEG and PMEA were active against herpes simplex virus, type 1 and 2 (HSV-1 and -2). (S)-HPMPA and (S)-cyclic HPMPA were active against varicella zoster virus (VZV). (S)-HPMPC was active against human cytomegalovirus (HCMV), a common cause of opportunistic infection in AIDS patients. (S)-HPMPA and (S)-cyclic HPMPA are active against adenovirus and vaccinia virus. PMEA, PMEDAP, and PMEMAP are active against human immunodeficiency virus (HIV), the human retrovirus responsible for AIDS. De Clercq, *et al.*, Antiviral Research, 8: 261-272 (1987).

Bronson, *et al.*, report on the series of acyclic nucleotide analogs having a common PME side chain attached to a purine or pyrimidine base which were prepared and selected for *in vivo* antiviral activity against retroviruses and herpes viruses. The adenine analog, PMEA, showed good *in vitro* activity against HIV and Rauscher murine leukemia virus (R-MuLV), and was more potent *in vivo* than 3'-azido-3'-deoxythymidine (AZT) in the treatment of R-MuLV in mice. PMEA also had a significant antiviral effect *in vivo* against murine cytomegalovirus (MCMV), and *in vitro* activity against HCMV. The guanine analog, PMEG, was exceptionally potent *in vitro* against herpes viruses. *In vivo*, PMEG was >50-fold more potent than acyclovir against HSV 1 infection in mice. Nucleotide Analogs as Antiviral Agents; ACS Symposium Series 401; Martin, J. C. Ed.: Washington, DC, 1989, Chapter 5, pp. 72-87. Kim, *et al.*, in J. Med. Chem., 33: 1207-1213 (1990), describe a similar series of compounds.

De Clercq, *et al.*, in Nature, 323: 464-467 (1986) state that (S)-HPMPA has potent and selective activity against a broad spectrum of DNA viruses, including HSV-1 and 2, VZV, thymidine kinase-deficient (TK<sup>-</sup>) mutants of herpes simplex HCMV, phocid herpesvirus type 1 (seal herpesvirus, SeHV), the simian herpesvirus platyrrhinae (HVP), suid herpesvirus type 1 (SHV-1, or pseudorabies virus or Aujeszky's disease virus), bovid herpesvirus type 1 (infectious bovine rhinotracheitis virus, BHV-1), equid herpesvirus type 1 (equine abortion virus, EHV-1), African swine fever (ASF) virus, vaccinia virus; and human adenoviruses, and retroviruses such as murine sarcoma virus (MSV). It is also reported that, in mice and rabbits *in vivo*, the compound is effective against both local and systemic infections with herpes simplex virus type 1, including herpetic keratitis caused by a TK<sup>-</sup> mutant which is resistant to the classical anti-herpes drugs.

European Patent Application 205,826, to De Clercq, *et al.*, published Dec. 30, 1986, discloses that HPMPA analogs are active against Moloney mouse sarcoma virus, and are expected to be effective against retroviruses in general. Reist and Sturm in PCT/U.S. 84/00737, published December 6, 1984 disclosed new phosphonic acid analogs of nucleoside phosphates which are useful as antivirals for incorporation into viral DNA.

Adenine phosphonic acid analogs and their synthesis are disclosed in the United Kingdom Patent application of Holy, *et al.*, GB 2,134,907A, published on August 22, 1984, and its related United States Patent, No. 4,659,825. A preferred example of one of these compounds, is known as (S)-9-((3-hydroxy-2-phosphonyl-methoxy)propyl)adenine (HPMPA). HPMPA was disclosed by E. DeClercq, *et al.*, in Nature, 323: 464-467, (1986), in Antiviral Research, 8: 261-272, (1987), and earlier by A. Holy, *et al.*, Nucleic Acids Research, Symposium Series No. 14: 277-278, (1984).

Phosphonyl-methoxyalkylpurine analogs have also been evaluated for their antitumor activity in murine tumor models. HPMPA, PMEA, and PMEG were found to be active against intraperitoneal P388 leukemia. PMEG was also found to be active against B16 melanoma. Rose, *et al.*, J. of the Nat. Cancer Inst., Vol. 82, No. 6 (1990).


A problem with nucleotides and other ionic organophosphate esters is their inability to traverse biological membranes. Liebman, *et al.*, J. Biol. Chem., 216: 823 (1955); Roll *et al.*, J. Biol. Chem., 220: 439 (1956). These compounds must, therefore, be given parenterally in order to achieve adequate serum levels to exert an antiviral effect.

Parenteral treatment is highly undesirable, especially with HIV infected patients. With HIV infected patients oral treatment is preferred since (i) HIV infected patients are very ill and need to be on chronic chemotherapy programs to maintain their health; (ii) the risk of using needle stick and presence of blood is high for health workers; (iii) disposal of infected needles is a problem; and (iv) the need for long-term maintenance therapy.


The inventors of this invention have carried out studies in order to circumvent the above-mentioned problem. The

present application, thus, relates to the preparation and use of a number of oral prodrugs of phosphonate nucleotide analogs.

In *J. Med. Chem.* 32:1457-1463 (1989), Bronson *et al.*, disclose the synthesis of HPMPC wherein the following compound is disclosed as an intermediate




In *Nucleotide Analogs as Antiviral Agents*, ACS Symposium Series 401, J.C. Martin, Ed., p. 72, American Chemical Society, Washington, D.C. (1989), Bronson *et al.*, disclose the synthesis of phosphonylmethoxy ether derivatives wherein the following compound was disclosed as an intermediate




wherein R is ethyl or isopropyl.

European Patent Application EP-270,885 of Webb, *et al.*, published June 15, 1988 discloses a process for the preparation of purin-9-ylalkyloxymethyl phosphonic acids, wherein several intermediates are produced in the practice of the process. One such intermediate is dialkylphosphonylmethyl which has the general structural formula




wherein R<sup>1</sup> and R<sup>2</sup>, independently, are selected from C<sub>1-6</sub> alkyl.

European Patent Application EP 253,412 of Holy, *et al.*, published January 20, 1988, discloses the preparation of a series of N-phosphonylmethoxyalkyl derivatives of pyrimidine and purine bases exhibiting antiviral activity, wherein in the practice of the process, several intermediates are produced. One such intermediate has the general structural formula



European Patent Application EP-269,947 of R. R. Webb, II, *et al.*, published on June 8, 1988, discloses a series antiviral agents which are phosphonomethoxyalkylene purine and pyrimidine derivatives having the following general structure




wherein  $\text{R}^3$  and  $\text{R}^4$  are independently selected from hydrogen,  $\text{C}_{1-16}$  alkyl, phenyl and phenyl- $\text{C}_{1-4}$ -alkylene.

The art compounds are generally distinguished from the compounds of the instant invention by the nature of the groups attached to the phosphorous atom. There is no disclosure or suggestion in the above references, or combination thereof, which would make obvious the use of a suitably protected phosphonate derivative prodrug for oral use.

This invention relates to prodrugs of phosphonate nucleotide analogs which exhibit antitumor activity and a broad spectrum of antiviral activity and some of which may be used orally.


The compounds of the instant invention comprise a diester-phosphonate link to nucleoside analogs of pyrimidine and purine bases. More particularly, it relates to compounds of the general structural formula as shown in Formula I



FORMULA I

wherein

B represents adenine (A), cytosine (C), guanine (G), thymine (T), uracil (U), 2,6-diamino purine (DAP), hypoxanthine (Hx),



wherein

Q is independently chosen from H, Cl,  $\text{NHR}^5$ ,  $\text{NR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH or  $\text{NCHN(R}^5)_2$ ;

R<sup>1</sup> and R<sup>2</sup> are identical or different and independently of one another are each OR<sup>4</sup>, NH<sub>2</sub>, NHR<sup>5</sup> or N(R<sup>5</sup>)<sub>2</sub>; R<sup>1</sup> and R<sup>2</sup> optionally being linked with each other to form a cyclic group, or R<sup>1</sup> or R<sup>2</sup> optionally being linked to R<sup>3</sup> to form a cyclic group;


R<sup>3</sup> represents C<sub>1</sub>-C<sub>20</sub> alkylene which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen; or R<sup>3</sup> is CH(CH<sub>2</sub>OR<sup>5</sup>)CH<sub>2</sub>, whereby R<sup>1</sup> and R<sup>2</sup> each independently may additionally represent OH, and R<sup>5</sup> is a hydrolyzable ester group;

R<sup>4</sup> represents a physiologically hydrolyzable ester group selected from CH<sub>2</sub>C(O)NR<sup>5</sup><sub>2</sub>, CH<sub>2</sub>C(O)OR<sup>5</sup>, CH<sub>2</sub>OC(O)R<sup>5</sup>, CH(R<sup>5</sup>)OC(O)R<sup>5</sup> (R, S, or RS stereochemistry), CH<sub>2</sub>C(R<sup>5</sup>)<sub>2</sub>CH<sub>2</sub>OH, or CH<sub>2</sub>OR<sup>5</sup>; or R<sup>4</sup> represents C<sub>4</sub>-C<sub>20</sub> alkyl, aryl-alkyl or aryl which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen provided that R<sup>1</sup> and R<sup>2</sup> are not simultaneously alkoxy;

R<sup>5</sup> represents C<sub>1</sub>-C<sub>20</sub> alkyl, aryl or aryl-alkyl which may be substituted or unsubstituted by substituents independently selected from the group consisting of hydroxy and halogen;

provided that those compounds of the above formula I are excluded wherein

B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil, R<sup>3</sup> is



wherein alk<sub>1</sub> is bonded to B, alk<sub>1</sub>, alk<sub>2</sub> and alk<sub>3</sub> are independently selected from a chemical bond or C<sub>1</sub>-C<sub>4</sub> alkylene, R<sup>a</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl and Q<sup>1</sup> is hydrogen or hydroxyl, and R<sup>1</sup> and R<sup>2</sup> are unsubstituted C<sub>4</sub>-C<sub>6</sub> alkoxy, phenoxy or phenyl-C<sub>1</sub>-C<sub>4</sub> alkoxy.

Included within the scope of the invention are the pharmaceutically acceptable acid addition salts, the metal salts and the solvate of the compounds of Formula I which may exist in various tautomeric forms.

In one aspect, the application relates to a process for the preparation of the compounds of Formula I.

In another aspect, the application relates to the use of the compounds of Formula I as a method for the treatment of viral infections in a mammal, which comprises administering an effective non-toxic dose of at least one compound of Formula I.


Another aspect of the application relates to the use of the compounds of Formula I as a method for inhibiting growth of a tumor in a mammal bearing a tumor which comprises administering an effective non-toxic dose of at least one compound of Formula I.

The compounds of Formula I are prodrugs of phosphonate nucleotides and have the same utility as the known or parent nucleotide analog. Thus the compounds of Formula I are useful as antiviral and antitumor agents.

The compounds of the present invention provide marked advantages over known nucleotides or analogs thereof in that these compounds are orally active.

The most preferred compounds of the invention are listed below, and experimental details for their preparation and characterization follow. Those which are not shown by specific example are readily prepared by analogous procedures.

A preferred example of the compounds of the instant invention are the compounds having the general structural formula as shown in Formula (II):




FORMULA I I

wherein

- B, R¹ and R² are as described in Formula I, provided that when Q is NCHN(R⁵)₂, then R⁵ is not CH₃;  
 X represents hydrogen, CH₂OR⁶ (R;S; or RS stereochemistry), or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is CH₂OR⁶, R¹ and R², may additionally be independently chosen from OH; and  
 R⁶ is a hydrolyzable ester group.

- Another preferred example of the compounds of the instant invention are the compounds having the general structural formula as shown in Formula (III):




FORMULA I I I

wherein

- B, and R¹ are as previously described in Formula I;  
 X represents hydrogen, CH₂OR⁶ (R;S; or R,S stereochemistry) or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is CH₂OR⁶, R¹ may additionally be OH; and  
 R⁶ is a hydrolyzable ester group;  
 R⁷ represents OH, NH₂, NHR⁵, or NR⁵₂; and  
 R⁵ is as described in Formula I, provided that those compounds of the above formula III are excluded wherein B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil, X is alk₂-Q¹, wherein alk₂ selected from a chemical bond or C₁-C₄ alkylene, Q¹ is hydrogen or hydroxyl, and R¹ and R⁷ are OH, unsubstituted C₄-C₆ alkoxy, phenoxy or phenyl-C₁-C₄ alkoxy.

- Still another preferred example of the compounds of the instant invention are the compounds having the general structural formula as shown in Formula (IV):




FORMULA I V

wherein

$R^8$  and  $R^9$  are identical or different and independently of one another are each  $NR^{12}$ , or oxygen;  
 $R^{10}$  and  $R^{11}$  are identical or different and independently of one another are each hydrogen, or  $R^5$ ;  
 $R^{12}$  represents hydrogen or a lower alkyl;  
 $m$  and  $n$  are identical or different and independently of one another are each 0 or 1;  
 $B$  and  $R^5$  are as described in Formula I; and  
 $X$  is as described in Formula II.

Yet another preferred example of the compounds of the instant invention are the compounds having the general structural formula as shown in Formula V



FORMULA V

\* stereochemistry is R, S, or RS

wherein

$R^{13}$  represents  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$ , or OH, provided that  $R^{13}$  is not OH when B is A or C; and  
 $B$ ,  $R^4$  and  $R^5$  are as described in Formula I.

The term "C<sub>1</sub> to C<sub>20</sub> alkyl" as used herein and in the claims (unless the context indicates otherwise) means saturated or unsaturated, branched or straight chain hydrocarbon group having 1 to 20 carbon atoms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, etc. Unless otherwise specified in the particular instance, the term "substituted or unsubstituted" as used herein and in the claims is intended to mean hydrocarbon group wherein an atom, element or group is regarded as having replaced a hydrogen atom, said substituted alkyl groups are preferably substituted with a member selected from the group consisting of hydroxy and halogen.

The term "prodrug" as used herein and in the claims (unless the context indicates otherwise) denotes a derivative of an active drug which is converted after administration back to the active drug. More particularly, it refers to derivatives of nucleotide phosphonates antiviral drugs which are capable of undergoing hydrolysis of the ester moiety or oxidative cleavage of the ester or amide moiety so as to release active, free drug. The physiologically hydrolyzable groups serve as prodrugs by being hydrolyzed in the body to yield the parent drug per se, and thus, the prodrugs of the present invention are preferably administered orally.

#### Synthesis of the phosphonate nucleotide analogs

The phosphonate nucleotide analogs are known compounds and therefore, the compounds as such and their chemical synthesis are not a part of the present invention. The synthesis of a number of phosphonate nucleotide analogs have been described in the literature.

For example, the synthesis of the phosphonates PME is disclosed in Holy and Rosenberg, Collect. Czech. Chem. Commun., 52:2801, (1987), and Bronson, et al, Nucleotide Analogues as Antiviral Agents, ACS Symposium Series 401, J.C. Martin, Ed., p. 72, American Chemical Society, Washington, D.C. (1989).

Bronson, et al, J. Med. Chem., 32: 1457-1463 (1989) discloses the preparation of HPMP from (R)-2,3-O-isopropylideneglycerol.


European Patent Application 253,412, published January 20, 1988 to Holy, et al, discloses methods for the preparation of PME and HPMP analogs of pyrimidine and purine bases.

Recently Holy et al Collect. Czech. Chem. Commun., 54: 2190-2210 (1989), described the preparation of N-(2-phosphorylmethoxy-ethyl) ("PME") analogs of purine and pyrimidine bases, as analogs of the antiviral 9-(2-phosphorylmethoxyethyl)adenine ("PMEA"). The synthesis consists of alkylation of alkali metal salts of heterocyclic bases or their N- or O-substituted analogs with diethyl 2-p-toluenesulfonyloxyethoxymethylphosphonate, 2-chloroethoxymethylphosphonate, or 2-bromoethoxymethylphosphonate. The obtained N-(2-diethoxyphosphonylmethoxyethyl) analogs of heterocyclic bases were treated with bromotrimethylsilane to give phosphonic acids. The phosphonic acids were prepared from pyrimidines (uracil, cytosine and their 5-methyl analogs), purines (adenine and its N<sup>6</sup> and C(2)-substituted analogs, hypoxanthine, guanine, 6-hydrazinopurine and 6-methylthiopurine etc.) and their analogs (3-deazaadenine etc.).


The synthesis of HPMPA is disclosed in Holy, Rosenberg, and Dvorakova, Collect. Czech. Chem. Commun. 54:2190 (1989).

#### Synthesis of dialkyl phosphonates


Quast, et al, Synthesis 490 (1974), has shown that dichlorophosphonates can be prepared by reacting phosphonates with  $\text{PCl}_5$ :




Moedritzer, K. CA 82, 86340, has shown that dichlorophosphonates can be prepared by reacting dimethylphosphonates with thionyl chloride.



Stowell, et al, (Tetrahedron Lett. 3261, (1990)) has shown that dichlorophosphonates can be reacted with alcohols or amines to give dialkylesters or dialkylamides:





The substituted phosphonates of the present invention were prepared by several methods: 1) Reaction of the phosphonate with thionyl chloride to give the dichlorophosphonate which was reacted further to give the disubstituted phosphonate:




2) Mono substituted phosphonates were obtained by the basic hydrolysis of the disubstituted phosphonate:






3) The monosubstituted phosphonates were chlorinated as before and reacted with a different alcohol or amine to give variably substituted phosphonates:



4) Diacyloxyalkyl phosphonates were obtained by reaction of the unsubstituted phosphonate with a substituted chloromethyl ether:



#### PROTOCOL FOR DETERMINING ORAL BIOAVAILABILITY OF PRODRUGS

Groups of rats, 3 rats per group were given a single iv dose of 30 mg/kg of PME A or a single oral dose of 30 mg-equiv/kg of PME A or PME A prodrug. Urine was collected in 0-24 hr and 24-48 hr intervals and analyzed for concentration of PME A. The bioavailability of PME A based on urinary excretion data and the bioavailability of PME A when given as a prodrug was determined. The results are summarized below.

ORAL BIOAVAILABILITY OF SELECTED PME A PRODRUGS IN RATS	
COMPOUND OF EXAM- PLE NO.	ABSOLUTE BIOAVAILA- BILITY
1 (PMEA)	7.8
8	17.0
11	15.4
12	14.6
13	34.9**
14	6.5
15	14.2
21	16.2
31	14.0
32	11.1

\*\*DETECTED AS THE MONOETHYL ESTER

IN VITRO ACTIVITY OF SELECTED PME A PRODRUGS AGAINST HSV-2 (G STRAIN)		
COMPOUND OF EXAM- PLE NO.	ID <sub>50</sub> (μg/mL) <sup>a</sup>	TOXICITY (μg/mL)
1 (PMEA)	39	>166
8	0.28	100
11	0.17	100
12	<0.1	100
13	3.3	100
14	8.1	100
15	>100	100
21	110	>166
31	42	>166
32	34	>166


<sup>a</sup>DOSE WHICH GIVES A 50 % REDUCTION OF PLACQUE FOR-  
MATION

The compounds of Formula I may be formulated for oral or parenteral use in a conventional manner using known pharmaceutical carriers and excipients, and they may be presented in unit dosage form or in multiple dose containers. The compositions may be in the form of tablets, capsules, solutions, suspensions or emulsions. These compounds may also be formulated as suppositories utilizing conventional suppository bases such as cocoa butter or other fatty materials. The compounds may, if desired, be administered in combination with other antiviral antibiotics.

When provided in unit dosage forms, the compositions may contain from about 0.1 to about 100 mg/kg/dose of the active ingredient of Formula I. The dosage of the compounds of Formula I is dependent on such factors as the weight

and age of the patient, as well as the particular nature and severity of the disease, and within the discretion of the physician. The dosage for adult human treatment may vary depending on the frequency and route of administration.

The following examples are intended for illustrative purpose only and are not to be construed as limiting the invention in sphere or scope. All temperatures are understood to be in degrees in C when not specified. The nuclear magnetic resonance (NMR) spectral characteristics refer to chemical shifts ( $\delta$ ) expressed in parts per million (ppm) versus tetramethylsilane (TMS) as reference standard. Except where otherwise noted,  $^1\text{H}$  spectra were recorded at 300 MHz and  $^{13}\text{C}$  spectra were recorded at 75 MHz. The relative area reported for the various shifts in the proton NMR spectral data corresponds to the number of hydrogen atoms of a particular functional type in the molecule. NMR assignments are based on the numbering system shown below:



The nature of the shifts as to multiplicity is reported as broad singlets (bs), singlets (s), multiplet (m), doublet (d), doublet of doublets (dd), triplet (t), or quartet (q). Coupling constants are given in hertz. When not specified, abbreviations employed are standard American Chemical Society (ACS) abbreviations as entered on the ACS Style Guide. The infrared (IR) spectral descriptions include only absorption wave numbers ( $\text{cm}^{-1}$ ) having functional group identification value. All compounds gave satisfactory elemental analyses, or high resolution mass spectrometry (HRMS).

#### I. GENERAL EXPERIMENTAL METHODS FOR COMPOUNDS LISTED IN TABLE I:

The compounds listed in Table I were synthesized by the corresponding method given at the end of the table. The reaction time, temperature and yield are given in Table I. The structure of the examples corresponds to either Figure 1 or Figure 2 given at the top of Table I. Spectral data for all compounds are given in the Examples which follow.

TABLE I. STRUCTURES AND EXPERIMENTAL DATA FOR PMEA PRODRUGS




FIGURE 1

FIGURE 2

STRUCTURE					EXPERIMENTAL			
Example No.	Fig	R1'	R2'	Q'	Method	Temp.°C	Time(h)	Yield %
2	1	iPrO	-R'	NH <sub>2</sub>	e			
3	1	iPrO	HO	OH	e			

STRUCTURE				EXPERIMENTAL				
Example No.	Fig	R <sup>1</sup>	R <sup>2</sup>	Q <sup>1</sup>	Method	Temp °C	Time (h)	Yield %
4	1	iPrO	-R <sup>1</sup>	Cl	e			
5	1	iPrO	-R <sup>1</sup>	H	e			
8	1	t-BuC(O)OCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	e	-	-	-
9	1	t-BuC(O)OCH <sub>2</sub> O	iPrO	NH <sub>2</sub>	e			
10	1	(CH <sub>3</sub> ) <sub>3</sub> N(+) (CH <sub>2</sub> ) <sub>2</sub> O	O <sup>-</sup>	NH <sub>2</sub>	e	-	-	-
11	1	EtC(O)OCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	e			
12	1	iPrC(O)OCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	e			
13	1	iPrC(O)OCH <sub>2</sub> O	CH <sub>3</sub> CH <sub>2</sub> O	NH <sub>2</sub>	e			
14	1	tBuC(O)OCH <sub>2</sub> O	OH	NH <sub>2</sub>	e			
15	1	iPrO	PhO	NH <sub>2</sub>	e			

EXPERIMENTAL				
Example No.	Fig	R <sub>1</sub> '	R <sub>2</sub> '	Q'
16	1	t-BuC(O)OCH <sub>2</sub> O	Et <sub>2</sub> NC(O)CH <sub>2</sub> O	NH <sub>2</sub>
17	2	O	H	NH <sub>2</sub>
18	1	Et <sub>2</sub> N	-R <sup>1</sup>	NH <sub>2</sub>
19	1	iPrO	O <sup>-</sup>	NH <sub>2</sub>
20	2	O	CH <sub>3</sub>	NH <sub>2</sub>
21	1	HO(CH <sub>2</sub> ) <sub>3</sub> O	O <sup>-</sup>	NH <sub>2</sub>
22	1	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>7</sub> O	HO	NH <sub>2</sub>
23	1	H <sub>2</sub> NCH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> NH-	HO	NH <sub>2</sub>
EXPERIMENTAL				
	Method	Temp °C	Time (h)	Yield %
	e			
	c	22	16	
	C	22	16	18
	B	60	24	75
	C	40	24	56
	B	60	2	50
	E	70	16	29
	B	60	1.5	61

STRUCTURE				EXPERIMENTAL				
Example No.	Fig	R1'	R2'	Q'	Method	Temp °C	Time (h)	Yield %
24	1	HOCH <sub>2</sub> C(CH <sub>3</sub> ) <sub>2</sub> CH <sub>2</sub> O	HO	NH <sub>2</sub>	D	60	1.5	80
25	2	NH	CH <sub>3</sub>	NH <sub>2</sub>	C	40	24	46
26	2	NCH <sub>3</sub>	H	NH <sub>2</sub>	F	82	24	27
27	1	Et <sub>2</sub> NC(O)CH <sub>2</sub> O	O <sup>-</sup>	NH <sub>2</sub>	F; D <sup>a,d</sup>	0; 22	20; 0.3	19
28	1	HOC(O)CH <sub>2</sub> O	HO	NH <sub>2</sub>	D	22	0.1	66
29	1	BuOC(O)CH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	F	82	1	44
30	1	EtOC(O)CH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	F	82	2	51
31	1	PhO	O <sup>-</sup>	NH <sub>2</sub>	D	22	1	76
32	1	PhO	-R <sup>1</sup>	NH <sub>2</sub>	F	22	20	38
33	1	iPr <sub>2</sub> NC(O)CH <sub>2</sub> O	O <sup>1-</sup>	NH <sub>2</sub>	F, D <sup>a,d</sup>	22; 22	0.8; 0.3	8

STRUCTURE				EXPERIMENTAL				
Example No.	Fig	R <sup>1</sup>	R <sup>2</sup>	Q <sup>1</sup>	Method	Temp °C	Time (h)	Yield %
34	1	pNO <sub>2</sub> PhCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	F	82	1	15
35	1	pNO <sub>2</sub> PhCH <sub>2</sub> O	O	NH <sub>2</sub>	D	60	20	78
36	1	CCl <sub>3</sub> CH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	F	82	20	44
37	1	CCl <sub>3</sub> CH <sub>2</sub> O	HO	NH <sub>2</sub>	D	60	1	69
38	1	PhC(O)OCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	e	22	20	9
39	1	pCF <sub>3</sub> PhCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	F	82	3	15
40	1	HOCH <sub>2</sub> CF <sub>2</sub> CH <sub>2</sub> O	HO	NH <sub>2</sub>	F; D <sup>a, d</sup>	82; 22	1; 1	18
41	1	pCF <sub>3</sub> PhCH <sub>2</sub> O	HO	NH <sub>2</sub>	D	60	20	82
42	1	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> NH	-R <sup>1</sup>	NH <sub>2</sub>	F	82	20	27
43	1	(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	A	60	3	75




STRUCTURE				EXPERIMENTAL				
Example No.	Fig	R <sup>1</sup>	R <sup>2</sup>	Q <sup>1</sup>	Method	Temp.°C	Time(h)	Yield %
44	1	(CH <sub>3</sub> ) <sub>2</sub> CH(CH <sub>2</sub> ) <sub>2</sub> O	-R <sup>1</sup>	NH <sub>2</sub>	A	80	1.5	77

- The crude product obtained from method F was employed directly in method D. Temperature and time are given for methods F and D, respectively.
- The impure product obtained from column chromatography was recrystallized from 25% MeOH/CH<sub>3</sub>CN.
- The impure product obtained from column chromatography was recrystallized from CH<sub>3</sub>CN.
- See detailed experimental section for synthesis of hydroxacetamides and difluoroalcohol.
- Where no method is given, see detailed experimental section for specific examples.

## METHOD OF SYNTHESIS FOR THE COMPOUNDS OF TABLE 1


A: A suspension of 1.00 g (3.66 mmol) of PMEA (1) in 50 mL of thionyl chloride was refluxed for 1 h (see eq. 1). The homogeneous, orange-red solution was cooled and the solvents were removed in vacuo to afford crude dichlorophosphonate 2. The dichloride was taken up in alcohol or amine 3 and stirred at the temperature and the time

given in Table I. After cooling the reaction to room temperature the solvents were removed in vacuo. The residue was purified on a 30 mm flash chromatography column, eluting with 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to afford 4. (See eq. 1)



(eq. 1)

B: An aqueous suspension of 4 was treated with 4 equivalents of NaOH for the time and temperature given in Table I (see eq. 2). The mixture was cooled to room temperature and acidified until pH 8. The majority of the solvent was evaporated and the residue was purified on a C-18 silica gel column, eluting with a gradient of 0-25% MeOH/H<sub>2</sub>O. The fractions containing the product were combined and evaporated to give 5. (See eq. 2)



(eq. 2)

C: This reaction was performed similarly to method A, except crude dichlorophosphate 2 was suspended in 30 mL of methylene chloride before adding alcohol or amine 3 (see equation 1).

D: This reaction was performed similarly to method B, except after cooling to room temperature, the reaction was acidified to pH 1.5. (See equation 2).

E: This reaction was run similarly to method B, except after cooling to room temperature the reaction was suspended in 20 mL of water. The mixture was acidified until pH approximately 3-4. The resulting solid was collected

and washed with water. The filtrate was cooled to 0 °C and the resulting solid was collected and washed with cold water. The solids were combined and dried overnight at 0.005 mm to afford 106 mg (0.23 mmol) of mono-octyl-PMEA.

- 5 F: This reaction was performed similarly to method A, except crude dichlorophosphonate 2 was suspended in 30 mL of acetonitrile before adding alcohol or amine 3 (see equation 1).

#### SPECIFIC EXPERIMENTAL METHODS FOR COMPOUNDS LISTED IN TABLE I.

##### 10 EXAMPLE 1

##### Synthesis of 9-(2-Phosphonylmethoxy)ethyladenine (PMEA).

A solution of PMEA diisopropyl ester (75.5 g, 0.21 mol) in 800 mL of anhydrous acetonitrile was treated with bromotrimethylsilane (258 g, 1.69 mol). The resulting clear, yellow solution was stirred at room temperature under argon for about 16 hours. The reaction mixture was concentrated in vacuo and the yellow residue was placed under high vacuum for about 5 hours. 400 mL of water was added next, causing immediate formation of a white precipitate. 500 mL of acetone was added and the pale yellow slurry was stirred at room temperature for about 14 hours. The solid was collected by filtration, washing twice with 150 mL of acetone and once with 150 mL of anhydrous ether. An additional portion of solid was collected from the filtrate to provide a total of 55.0 g (90%) of PMEA as an off-white crystalline solid.

m.p. > 250°C; UV<sub>max</sub> (H<sub>2</sub>O) 208 nm ( $\epsilon$  = 19,600) 260 nm ( $\epsilon$  = 14,100); UV<sub>max</sub> (0.1 N HCl) 210 nm ( $\epsilon$  = 19,000) 260 nm ( $\epsilon$  = 13,700); UV<sub>max</sub> (0.1 N NaOH) 216 nm ( $\epsilon$  = 9,600) 262 nm ( $\epsilon$  = 14,500); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  8.14 (s, 1 H), 8.13 (s, 1 H), 7.27 (br s, 2 H, NH<sub>2</sub>), 4.32 (t, *J* = 5, 2 H, H-1'), 3.87 (t, *J* = 5, 2 H, H-2'), 3.59 (d, *J* = 9, 2 H, H-4'); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>)  $\delta$  151.10 (C-6), 148.70 (C-2), 146.28 (C-4), 143.80 (C-8), 118.05 (C-5), 69.94 (d, *J* = 10, C-2'), 66.27 (d, *J* = 160, C-4'), 43.15 (C-1').

##### EXAMPLE 2

##### 30 Synthesis of PMEA, di-(isopropyl ester).

A slurry of adenine (21.2 g, 157 mmol), 2-[(diisopropylphosphonyl)methoxy]ethyl methanesulfonate (50.0 g, 157 mmol, prepared according to the procedure described by J.J. Bronson *et al.* in *J. Med. Chem.*, 32: 1457, (1989)), and cesium carbonate (56.0 g, 173 mmol) in 160 mL of anhydrous DMF was heated to 120°C in a 3-necked, 500-mL, round-bottomed flask equipped with a mechanical stirrer and argon inlet adapter. The reaction mixture was stirred at 120°C for about 5 hours and then was allowed to cool to room temperature. Insoluble material was removed by filtration and the filtrate was concentrated in vacuo to give 66 g of a yellow solid. Purification by column chromatography on silica gel (10:1, elute with 3% to 5% to 7% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) provided 33 g of an off-white solid. Recrystallization from ethyl acetate provided 30.1 g (54%) of PMEA, diisopropyl ester as a white solid.

Mp 136-138°C; UV<sub>max</sub> (MeOH) 262 nm ( $\epsilon$  = 14,360); <sup>1</sup>H NMR (DMSO-*d*<sub>6</sub>)  $\delta$  8.15 (s, 1H), 8.09 (s, 1H), 7.21 (br s, exch, 2H, NH<sub>2</sub>), 4.50 (apparent octet, *J* = 6.5 Hz, 2H, 2POCH), 4.34 (t, *J* = 5 Hz, 2H, NCH<sub>2</sub>), 3.91 (t, *J* = 5 Hz, 2H, CH<sub>2</sub>OCH<sub>2</sub>P), 3.79 (d, *J* = 8 Hz, 2H, OCH<sub>2</sub>P), 1.18 (d, *J* = 6.5 Hz, 6H, POCH(CH<sub>3</sub>)<sub>2</sub>), and 1.13 (d, *J* = 6.5 Hz, 6H, POCH(CH<sub>3</sub>)<sub>2</sub>); <sup>13</sup>C NMR (DMSO-*d*<sub>6</sub>)  $\delta$  155.86 (C-6), 152.23 (C-2), 149.46 (C-4), 140.90 (C-8), 118.57 (C-5), 70.22 (d, *J* = 10 Hz, POCH), 70.05 (d, *J* = 12 Hz, CH<sub>2</sub>OCH<sub>2</sub>P), 64.50 (d, *J* = 165 Hz, OCH<sub>2</sub>P), 42.35 (NCH<sub>2</sub>), 23.61 [d, *J* = 7 Hz, POCH(CH<sub>3</sub>)<sub>2</sub>], and 23.52 [d, *J* = 7 Hz, POCH(CH<sub>3</sub>)<sub>2</sub>]; mass spectrum (methane DCI), *m/z* (rel intensity) 358 (MH<sup>+</sup>, 100), 344 (10), 316 (10).

50

Anal. Calc. for C <sub>14</sub> H <sub>24</sub> N <sub>5</sub> O <sub>4</sub> P:	C, 47.06;	H, 6.77;	N, 19.60.
Found:	C, 47.06;	H, 7.04;	N, 19.65.

55

## EXAMPLE 3

Synthesis of 9-(2-Phosphonylmethoxy)ethylhypoxanthine (PMEHx), mono-(isopropyl ester).

5 A solution of 6-chloro-9-(2-phosphonylmethoxy) ethylpurine, diisopropyl ester (1g, 2.65 mmol) in 27 mL of 1 N NaOH was heated at reflux for 1 h, cooled to room temperature, acidified to pH 1 with 1 N HCl and concentrated in vacuo. The residue was purified by C-18 silica gel column chromatography, eluting with 20% MeOH/H<sub>2</sub>O to afford 0.51 g (68%) of the title compound.

10 Mp 192-194°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 12.27 (1H, br s, NH), 8.04, 8.02 (2H, 2s, H-2, H-8), 4.39 (1H, septet, J=6, CH(CH<sub>3</sub>)<sub>2</sub>), 4.30 (2H, t, J=5, H-1'), 3.85 (2H, t, J=5, H-2'), 3.65 (2H, d, J=8.5, H-4'), 1.10 (6H, d, J=6, CH<sub>3</sub>). <sup>13</sup>C NMR (D<sub>2</sub>O) 157.57 (C=O), 149.94, 149.72 (C-2, C-4), 143.02 (C-8), 119.94 (C-5), 72.09 (CH(CH<sub>3</sub>)<sub>2</sub>, d, J=6), 71.70 (C-2', d, J=13), 67.80 (C-4', d, J=160), 47.05 (C-1), 25.28 (CH<sub>3</sub>, d, J=4), IR (KBr) 3423, 2979, 1716, 1642, 1587, 1562. MS (FAB) 317 (M+H, 100).

15

20

Anal. Calcd for C <sub>11</sub> H <sub>17</sub> N <sub>4</sub> O <sub>5</sub> P • 0.4 H <sub>2</sub> O:	C, 40.95;	H, 5.53;	N, 17.37.
Found:	C, 41.19;	H, 5.68;	N, 17.51.

## EXAMPLE 4

Synthesis of 6-Chloro-9-(2-Phosphonylmethoxy) ethylpurine, di-(isopropyl ester).

25

To a rapidly stirred solution of 9.86 g (63.8 mmol) of 6-chloropurine in 350 mL of anhydrous DMF was added 1.91 g (63.8 mmol) of sodium hydride (80% in mineral oil). The heterogeneous mixture was heated at 95 °C for about 20 hours, cooled to room temperature and concentrated in vacuo. The residue was purified by silica gel chromatography, eluting with 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 4.53 g of the title compound.

30

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.76 (1H, s, H-8), 8.63 (1H, s, H-2), 4.82 (2H, t, J=5, H-1'), 4.42 (2H, septet, J=6, CH(CH<sub>3</sub>)<sub>2</sub>), 3.93 (2H, t, J=5, H-2'), 3.75 (2H, d, J=8, H-4'), 1.11 (6H, d, J=6, CH<sub>3</sub>), 1.05 (6H, d, J=6, CH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 152.44 (C-6), 151.88 (C-2), 149.39 (C-4), 148.13 (C-8), 131.13 (C-5), 70.24 (CH(CH<sub>3</sub>)<sub>2</sub>, d, J=6), 70.00 (C-2', d, J=11), 64.64 (C-4', d, J=165), 43.43 (C-1'), 23.65 (CH<sub>3</sub>, d, J=4.5), 23.47 (CH<sub>3</sub>, d, J=4.5). IR (KBr) 3459, 3077, 2982, 2936, 1564. MS (methane/DCI) 377 (M+H, 100).

35

40

Anal. Calcd for C <sub>14</sub> H <sub>22</sub> N <sub>4</sub> O <sub>4</sub> Cl <sub>1</sub> P <sub>1</sub> :	C, 44.63;	H, 5.89;	N, 14.87.
Found:	C, 44.40;	H, 5.93;	N, 14.53.

## EXAMPLE 5

Synthesis of 9-(2-phosphonylmethoxy)ethylpurine, di-(isopropyl ester).

45

A solution of 6-Chloro-9-(2-phosphonylmethoxy) ethylpurine, diisopropyl ester (0.94 g, 2.5 mmol) in 20 mL of ethanol/cyclohexene (1:1) was treated with 0.5 g of Pd(OH)<sub>2</sub>/C. The reaction was stirred at reflux for about 20 hours, diluted with hot ethanol and filtered through celite. The filtrate was concentrated in vacuo and the residue was purified by flash column chromatography, eluting with 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to afford 0.49 g (58%) of the title purine as a clear yellow oil.

50

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 9.14, 8.92, 8.55 (3H, 3s, H-2, H-6, H-8), 4.47 (2H, t, J=5, H-1'), 4.42 (2H, septet, J=6, CH(CH<sub>3</sub>)<sub>2</sub>), 3.94 (2H, t, J=5, H-2'), 3.77 (2H, d, J=8, H-4'), 1.12 (6H, d, J=6, CH<sub>3</sub>), 1.05 (6H, d, J=6, CH<sub>3</sub>). IR 3459, 2982, 2937, 1597, 1581, 1506. MS (methane/DCI) 343 (M+H, 100), 329 (12), 301 (50).

55

Anal. Calcd for $C_{14}H_{23}N_4O_4P \cdot 0.25 H_2O$ :	C, 48.50;	H, 6.83;	N, 16.16.
Found:	C, 48.55;	H, 6.71;	N, 15.88.

## EXAMPLE 6

## Synthesis of hydroxyacetamides necessary for preparation of Example 20 and Example 24.

## (a) 2-hydroxy-N,N-diethylacetamide

A solution of 10.5 g (0.0702 mol) of 2-chloro-N,N-diethylacetamide in 35 mL of glacial acetic acid was refluxed for about 16 hours. The solvents were removed in vacuo, the last traces of acetic acid being azeotropically removed with toluene. The residue was dissolved in 125 mL of methanol and treated with 10.85 g (0.20 mol) of sodium methoxide. The reaction was stirred for about 3 hours and neutralized with Dowex 50X8-200 acidic ion exchange resin. The solvents were removed in vacuo and the residue was purified on a flash chromatography column, eluting with hexane/ethyl acetate 1:1 to give 6.75 g (73%) of 2-hydroxy-N,N-diethylacetamide.

## (b) 2-hydroxy-N,N-diisopropylacetamide

To a solution of 44.5 g (0.44 mol) of N,N-diisopropyl amine in 125 mL of hexane cooled to -78 °C was added dropwise 17.6 mL (0.22 mol) of chloroacetyl chloride. After completion of the addition, the cooling bath was removed and stirring was continued for about 30 minutes. The isopropylammonium chloride was removed by filtration through celite and the filtrate was stripped to give 30.5 g (77%) of 2-chloro-N,N-diisopropylacetamide. Hydrolysis of this compound as described above afforded a 45% yield of 2-hydroxy-N,N-diisopropylacetamide.

## EXAMPLE 7

## Synthesis of the difluoroalcohol necessary for the preparation of Example 31.

## (a) 2,2-Difluoro-3-hydroxy-propan-1-ol

A solution of 9.07 g (0.0521 mol) of 1,3-diacetyl acetone in 20 mL of DAST was stirred at 22 °C for 2 days, diluted with ethyl acetate, washed with saturated  $NaHCO_3$  and water, then dried over  $Na_2SO_4$  and concentrated to yield 9.54 g of 1,3-diacetyl-2,2-difluoropropane. The diacetyl-difluoropropane (7.53 g, 38.4 mmol) was dissolved in 300 mL of methanol and treated with 6.45 g (119 mmol) of sodium methoxide. After stirring at 22 °C for about 2.5 hours, the reaction was neutralized with Dowex 50X8-200 acidic ion exchange resin, filtered and stripped to give 3.7 g (86%) of the title compound.

## EXAMPLE 8

## Synthesis of PME A, di-(pivaloyloxymethyl ester)

To a rapidly stirred solution of 1.00 g (3.66 mmol) of PME A in 15 ml of anhydrous DMF was added 2.08 g (7.32 mmol) of N,N'-dicyclohexyl-4-morpholine carboxamidine and 2.75 g (18.33 mmol) of chloromethyl pivalate. The heterogeneous mixture became homogeneous after about 15 minutes and was then allowed to stir at 22°C for about 36 hours. The insolubles were filtered off and the filtrate was concentrated in vacuo. The residue was then partitioned between (50 ml) water and (50 ml) toluene, separated and the water layer was then extracted with (2 x 50 ml) toluene. The toluene layers were combined and concentrated in vacuo. The residue was purified by silica gel chromatography, eluting with 5% MeOH/ $CH_2Cl_2$  to give 0.59 g (32%) of the title compound.

$^1H$  NMR( $CDCl_3$ ) 8.32(1H, s, H-8), 7.91(1H, s, H-2), 5.77(2H, s,  $NH_2$ ), 5.63(4H, m,  $CH_2OP$ ), 4.37(2H, t, J=5.0, H-1'), 3.92(2H, t, J=5.0, H-2'), 3.82(2H, d, J=7.7, H-4'), 1.18(18H, s,  $CH_3$ ).  $^{13}C$  NMR ( $CDCl_3$ ) 177.55(C=O), 156.23(C-6), 153.45(C-2), 150.48(C-4), 142.05(C-8), 119.85(C-5), 82.04 ( $CH_2OP$ , d, J=6.0), 71.70(C-2', d, J=9.8), 65.86(C-4', d, J=167), 43.63(C-1'), 38.95(C=O), 27.11( $CH_3$ ). IR(KBr) 3366, 3178, 2976, 1754, 1660, 1600. MS(Isobutane/DCI) 502(M+H, 100).

Anal. Calcd. for $C_{20}H_{32}N_5O_8P_1$ :	C, 47.90;	H, 6.43;	N, 13.96.
Found:	C, 48.02;	H, 6.27;	N, 13.63.

## EXAMPLE 9

10 Synthesis of PMEA, (mono-isopropyl, mono-pivaloyloxymethyl) ester

To a rapidly stirred solution of 200 mg (0.6 mmol) of monoisopropyl PMEA (example 11) in 5 ml of anhydrous DMF was added 0.83 ml (6.0 mmol) of  $Et_3N$  and 0.45 g (3.0 mmol) of chloromethylpivalate. The heterogeneous mixture became homogeneous after addition of  $Et_3N$  and was then allowed to stir at 22°C for about 3 days. The mixture was concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 10% MeOH/ $CH_2Cl_2$  to give 190 mg (74%) of the title compound.

$^1H$  NMR( $CDCl_3$ ) 8.30(1H, s, H-8), 7.91(1H, s, H-2), 5.94(2H, s,  $NH_2$ ), 5.57(2H, d,  $J=12.5$ ,  $CH_2OP$ ), 4.73(1H, septet,  $J=6.2$ , CH), 4.36(2H, t,  $J=5.0$ , H-1'), 3.90(2H, t,  $J=5.0$ , H-2'), 3.75(2H, d,  $J=8.0$ , H-4'), 1.25(6H, d,  $J=6.2$ ,  $CH_3$ ), 1.17(9H, s,  $CH_3$ ).  $^{13}C$  NMR ( $CDCl_3$ ) 177(C=O), 155.51(C-6), 152.91(C-2), 149.8 (C-4), 141.43(C-8), 119.36(C-5), 81.90( $CH_2OP$ , d,  $J=5.6$ ), 72.18( $CHOP$ , d,  $J=7.0$ ), 71.19(C-2', d,  $J=10.0$ ), 65.78(C-4', d,  $J=16.7$ ), 43.37 (C-1'), 38.68( $(CH_3)_3C$ ), 26.84( $(CH_3)_3C$ ), 23.92( $CH_3CH$ , d,  $J=7$ ), 23.85( $CH_3CH$ , d,  $J=7$ ). IR(KBr) 3432, 1754, 1668, 1602. MS(FAB) 430(M+H, 100).

Anal. Calcd. for $C_{17}H_{28}N_5O_6P_1 \cdot 0.50 H_2O$ :	C, 46.56;	H, 6.66;	N, 15.98.
Found:	C, 46.50;	H, 6.61;	N, 15.81.

## EXAMPLE 10

Synthesis of PMEA, mono-(choline ester)

A suspension of 2.00 g (7.33 mmol) of PMEA in 30 ml of thionyl chloride was refluxed for about 1 hour. The homogeneous, orange-red solution was cooled and the solvents were removed in vacuo to afford crude dichlorophosphate. The dichloride was taken up in 40 ml of acetonitrile and then treated with 2.00 g (32.34 mmol) of anhydrous ethylene glycol at reflux for about 16 hours. After cooling to 22°C, the solvents were removed in vacuo. The residue was purified by silica gel chromatography, eluting with MeOH/ $CH_2Cl_2$ / $NH_4OH$  30/70/1 to give 1.42 g (65%) of mono(chloroethyl)ester.

A suspension of 460 mg (1.37 mmol) of the above compound in 30 ml of MeOH was saturated with  $Me_3N$  gas at 0°C. The reaction mixture was then sealed in a metal bomb and heated at 65°C for about 2 days.

After cooling the reaction to 22°C, the solvents were removed in vacuo and the residue was purified by C-18 chromatography, eluting with 15% MeOH/ $H_2O$  to give 270 mg (35% from PMEA) of the title compound.

$^1H$  NMR( $CD_3OD$ ) 8.24(1H, s, H-8), 8.20(1H, s, H-2), 4.42(2H, t,  $J=5.0$ , H-1'), 4.12(2H,  $CH_2CH_2OP$ ), 3.89(2H, t,  $J=5.0$ , H-2'), 3.64(2H, d,  $J=9.0$ , H-4'), 3.47 (2H, m,  $CH_2OP$ ), 3.14(9H, s,  $CH_3$ ).  $^{13}C$  NMR ( $CD_3OD$ ) 157.55(C-6), 154.03(C-2), 151.02(C-4), 144.02(C-8), 120.15(C-5), 72.04(C-2'), 68.24(C-4', d,  $J=15.9$ ), 68.05 ( $CH_2OP$ ), 60.10( $CH_2CH_2OP$ , d,  $J=4.9$ ), 55.02( $CH_3$ ), 54.98( $CH_3$ ), 54.92( $CH_3$ ), 44.95(C-1'). IR(KBr) 3396, 1648, 1602, 1480. MS(FAB) 359(M+H, 30).

Anal. Calcd. for $C_{13}H_{23}N_5O_4P_1 \cdot 2.5H_2O$ :	C, 38.60;	H, 7.00;	N, 20.78.
Found:	C, 38.26;	H, 6.60;	N, 20.47.

## EXAMPLE 11

Synthesis of PME A, di-(propionyloxymethyl ester)

To a rapidly stirred solution of 1.00 g (3.66 mmol) of PME A in 15 ml of anhydrous DMF was added 2.08 g (7.32 mmol) of N,N'-dicyclohexyl-4-morpholine carboxamidine and 2.23 g (18.3 mmol) of chloromethylpropionate. The heterogeneous mixture became homogeneous within 30 minutes and was then allowed to stir at 22°C for about 5 days. The insolubles were filtered off and the filtrate was concentrated in vacuo. The residue was purified twice by silica gel chromatography (200:1), eluting with 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 0.14g (9%) of the title compound.

<sup>1</sup>H NMR(CDCl<sub>3</sub>) 8.29(1H, s, H-8), 7.88(1H, s, H-2), 5.65(2H, s, NH<sub>2</sub>), 5.60(4H, m, CH<sub>2</sub>OP), 4.35(2H, t, J=5.0, H-1), 3.89(2H, t, J=5.0, H-2'), 3.80(2H, d, J=7.8, H-4'), 2.34(4H, q, J=7.5 CH<sub>3</sub>CH<sub>2</sub>), 1.10 (6H, t, J=7.5, CH<sub>3</sub>). IR(KBr) 3290, 3122, 1766, 1666, 1602. MS(FAB) 446(M+H, 100).

Anal. Calcd. for C <sub>16</sub> H <sub>24</sub> N <sub>5</sub> O <sub>6</sub> P <sub>1</sub> :	C, 43.15;	H, 5.43;	N, 15.72.
Found:	C, 43.07;	H, 5.46;	N, 15.42.

## EXAMPLE 12

Synthesis of PME A, di-(isobutyroxymethyl ester)

To a rapidly stirred solution of 1.00 g (3.66 mmol) of PME A in 15 ml of anhydrous DMF was added 2.08 g (7.32 mmol) of N,N'-dicyclohexyl-4-morpholine carboxamidine and 2.48 g (18.3 mmol) of chloromethylisobutyrate. The heterogeneous mixture became homogeneous within 30 minutes and was then allowed to stir at 22°C for 5 days. The mixture was concentrated in vacuo, partitioned between (50 ml) water and (50 ml) toluene. The aqueous layer was extracted with (250 ml) toluene and the combined organic layer was concentrated in vacuo. The residue was purified by silica gel chromatography, eluting with 5% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 0.16 g (9%) of the title compound.

<sup>1</sup>H NMR(CDCl<sub>3</sub>) 8.31(1H, s, H-8), 8.28(1H, s, H-2), 5.68(2H, s, NH<sub>2</sub>), 5.59(4H, m, CH<sub>2</sub>OP), 4.33(2H, t, J=5.0, H-1), 3.88(2H, t, J=5.0, H-2'), 3.78(2H, d, J=7.7H, H-4'), 2.52(2H, apparent heptet, J=7.0, CH), 1.11(6H, d, J=7.0, CH<sub>3</sub>). IR(KBr) 3360, 2980, 1758, 1660, 1602. MS(Isobutane/DCI) 474(M+H, 100).

Anal. Calcd. for C <sub>18</sub> H <sub>28</sub> N <sub>5</sub> O <sub>6</sub> P <sub>1</sub> · 0.65 H <sub>2</sub> O:	C, 44.56;	H, 6.09;	N, 14.44.
Found:	C, 45.67;	H, 5.96;	N, 14.79.

## EXAMPLE 13 (for comparison)

Synthesis of PME A, (mono-ethyl, mono-isobutyryloxymethyl) ester

To a rapidly stirred solution of 400 mg (1.33 mmol) of monoethyl PME A in 15 ml of anhydrous DMF was added 2.00 ml (14.3 mmol) of Et<sub>3</sub>N and 1.0 g (6.7 mmol) of chloromethylpivalate. The heterogeneous mixture became homogeneous after addition of Et<sub>3</sub>N and was then allowed to stir at 22°C for 2 days. The mixture was concentrated in vacuo and the residue was purified by silica gel chromatography, eluting with 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 180 mg (33%) of the title compound.

<sup>1</sup>H NMR(CDCl<sub>3</sub>) 8.32(1H, s, H-8), 7.92(1H, s, H-2), 5.74(2H, s, NH<sub>2</sub>), 5.62(2H, m, OCH<sub>2</sub>OP), 4.38 (2H, t, J=5.0, H-1), 4.10(2H, m, CH<sub>3</sub>CH<sub>2</sub>OP), 3.92(2H, t, J=5.0, H-2'), 3.79(2H, d, J=8.0, H-4'), 1.27 (3H, t, J=7.0, CH<sub>3</sub>CH<sub>2</sub>), 1.18 (9H, s, ((CH<sub>3</sub>)<sub>3</sub>C)). <sup>13</sup>C NMR (CDCl<sub>3</sub>) 176.87(C=O), 155.40(C-6), 152.94(C-2), 149.8(C-4), 141.51(C-8), 119.7(C-5), 81.85(CH<sub>2</sub>OP, d, J=6.2), 71.26(C-2', d, J=10.2), 65.46(C-4', d, J=16.7), 62.73(CH<sub>2</sub>CH<sub>3</sub>, d, J=7.0), 43.49(C-1'), 38.70((CH<sub>3</sub>)<sub>3</sub>C), 26.84((CH<sub>3</sub>)<sub>3</sub>C), 16.27(CH<sub>2</sub>CH<sub>3</sub>, d, J=5.8). IR(KBr) 3288, 3120, 2982, 1752, 1666, 1600. MS(FAB) 416(M+H, 100).



Anal. Calcd. for $C_{16}H_{25}N_5O_5P_1 \cdot 0.5H_2O$ :	C, 45.28;	H, 6.41;	N, 16.51.
Found:	C, 45.47;	H, 6.34;	N, 16.55.

## EXAMPLE 14

10 Synthesis of PMEA, mono-(pivaloyloxymethyl ester)

To a solution of sodium hydride (0.95 g, 80%, 31.7 mmol) and benzylalcohol (6.8 ml, 63.5 mmol) in anhydrous DMSO (50 ml) was added with stirring a solution of PMEA, diphenyl ester (3.4 g, 8 mmol, example 26) in DMSO (50 ml). The mixture was allowed to stir at 22°C for 1 h and concentrated to a volume of approximately 25 ml. EtOAc (200 mL) was added and the precipitate was collected by vacuum filtration. The precipitate was purified by C-18 chromatography, eluting with 20% MeOH/H<sub>2</sub>O to give 2.09 g (68%) of PMEA, monobenzylester, sodium salt.

To 600 mg (1.56 mmoles) of the above compound in 14 ml of anhydrous DMF was added 2.16 ml (15.5 mmoles) of Et<sub>3</sub>N and 1.44 g (9.61 mmol) of chloromethylpivalate. The mixture was allowed to stir at 22°C for 2 days, concentrated in vacuo and the resulting residue was used crude in the following step.

To a stirred solution of the crude mixed ester (300 mg) in 17 ml of EtOH and 17 ml of H<sub>2</sub>O was added 3.45 ml of cyclohexene and 0.225g of 20% Pd(OH)<sub>2</sub>/C. The mixture was heated at reflux for 1h, concentrated in vacuo and the residue purified by C-18 chromatography, eluting with 100% H<sub>2</sub>O to give 270 mg (31% from PMEA, diphenyl ester) of the title compound.

<sup>1</sup>H NMR(d<sub>6</sub>-DMSO) 8.09(2H, s, H-8, H-2), 7.17(2H, s, NH<sub>2</sub>), 5.44(2H, m, CH<sub>2</sub>OP), 4.26(2H, t, J=5.0, H-1'), 3.83(2H, t, J=5.0, H-2'), 3.47(2H, d, J=8.0, H-4'), 1.04(9H, s, CH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 176.70(C=O), 155.98(C-6), 152.39(C-2), 149.55(C-4), 141.30(C-8), 118.59(C-5), 83.14(CH<sub>2</sub>OP), 69.89(C-2'), 64.5(C-4'), 42.84 (C-1'), 38.13 ((CH<sub>3</sub>)<sub>3</sub>C) 26.69(CH<sub>3</sub>). IR(KBr) 3360, 1742, 1648, 1602. MS(FAB) 386(M-H, 100). HRMS:

Calculated:	388.1386.
Found:	388.1377.

## EXAMPLE 15

40 Synthesis of PMEA, (mono-isopropyl, mono-phenyl) ester

A suspension of 0.75 g (2.1 mmol) of monophenyl PMEA in 20 ml of thionyl chloride was refluxed for 1 h. The homogeneous, orange-red solution was cooled and the solvents were removed in vacuo to afford crude monochlorophosphonate. The residue was taken up in 40 ml of isopropyl alcohol and stirred for 16 h at 22°C. The solvents were removed in vacuo and the residue was purified by silica gel chromatography, eluting with 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 0.24g (29%) of the title compound.

Mp 96-99°C. <sup>1</sup>H NMR(CDCl<sub>3</sub>) 8.31(1H, s, H-8), 7.87(1H, s, H-2), 7.19(5H, m, Ph), 5.96(2H, s, NH<sub>2</sub>), 4.80 (1H, apparent heptet, J=6.2, CH), 4.36(2H, t, J=5.0, H-1'), 3.93(2H, t, J=5.0, H-2'), 3.86(2H, d, J=7.9, H-4'), 1.26(3H, d, J=6.2, CH<sub>3</sub>), 1.21(3H, d, J=6.2 CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>) 155.52(C-6), 152.88(C-2), 150.13(ArC, d, J=8.3), 149.89(C-4), 141.46(C-8), 129.71(ArC), 125.14(ArC), 120.50(ArC, d, J=4.5), 119.43(C-5), 72.65(CH, d, J=7.3), 71.18(C-2', d, J=10.6), 65.27(C-4', d, J=167.5), 43.45(C-1'), 23.93(CH<sub>3</sub>, d, J=4.5), 23.82(CH<sub>3</sub>, d, J=4.5). IR(KBr) 3290, 3116, 1670, 1600. MS(Isobutane/DCI) 392(M+H, 100).

Anal. Calcd. for $C_{17}H_{22}N_5O_4P_1$ :	C, 52.17;	H, 5.66;	N, 17.89.
Found:	C, 52.01;	H, 5.57;	N, 17.64.

## EXAMPLE 16

Synthesis of PMEA, (mono-N,N-diethylacetamide, mono-pivaloyloxymethyl) ester

To a suspension of 0.100 g (0.239 mmol) of PMEA, mono- N,N-diethylacetamide ester (sodium salt) (Example 24) in 2.5 mL of CH<sub>3</sub>CN was added 0.25 mL of Et<sub>3</sub>N, whereupon the reaction became homogeneous. To this mixture was added 0.17 mL (1.19 mmol) of chloromethyl pivalate. The reaction was stirred at 22°C for 24h, evaporated to dryness in vacuo, and purified on a 20 mm flash column. The title compound eluted with 10 % MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give 25 mg (21%) of a colorless oil.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) 8.25 (1H, s, H-8), 7.94 (1H, s, H-2), 6.26 (2H, s, NH<sub>2</sub>), 5.65 (1H, dd, J=12.3, 5.4, OCH<sub>2</sub>O), 5.60 (1H, dd, J=12.3, 4.8, OCH<sub>2</sub>O), 4.75 (1H, dd, J=14.7, 10.8, OCH<sub>2</sub>C(O)), 4.56 (1H, dd, J=14.5, 14.3, OCH<sub>2</sub>C(O)), 4.32 (2H, dd, J=5.7, 4.4, H-1'), 3.97 (2H, d, J=8.4, H-4'), 3.91 (2H, t, J=4.8, H-2'), 3.28 (2H, q, J=7.5, CH<sub>2</sub>CH<sub>3</sub>), 3.09 (1H, q, J=7.2, CH<sub>2</sub>CH<sub>3</sub>), 1.12 (9H, s, (CH<sub>3</sub>)<sub>3</sub>), 1.07 (3H, m, CH<sub>3</sub>CH<sub>2</sub>), 1.05, (3H, t, J=6.9, CH<sub>3</sub>CH<sub>2</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>) 177.85 (C(O)O), 166.25 (C(O)N), 156.34 (C-6), 153.48 (C-2), 150.49 (C-4), 142.22 (C-8), 119.79 (C-5), 81.94 ((CH<sub>3</sub>)<sub>3</sub>C), 81.71 (OCH<sub>2</sub>O), 71.55 (C-2', d, J=10), 65.10 (C-4', d, J=165), 63.99 (CCH<sub>2</sub>OP), 43.53 (C-1'), 41.03 (NCH<sub>2</sub>), 40.78 (NCH<sub>2</sub>), 27.00 ((CH<sub>3</sub>)<sub>3</sub>), 14.21 (CH<sub>3</sub>CH<sub>2</sub>), 13.00 (CH<sub>3</sub>CH<sub>2</sub>). MS (FAB) 501 (M+H, 100). IR 3500-3000, 2978, 1750, 1654, 1600, 1480, 1250.

Anal. Calcd for: C <sub>20</sub> H <sub>33</sub> N <sub>5</sub> O <sub>7</sub> P · 0.5 H <sub>2</sub> O	C, 47.15;	H, 6.72;	N, 16.50.
Found:	C, 47.30;	H, 6.58;	N, 16.14.

The following examples were prepared by the methods given in Table I.

## EXAMPLE 17

PMEA, cyclic propanyl diester

Mp 195-199°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.13 (1H, s, H-8), 8.12 (1H, s, H-2), 4.35 (2H, t, J=4.8, H-1'), 4.2 (4H, m, CH<sub>2</sub>OP), 3.95 (2H, d, J=8.8, H-4'), 3.86 (2H, t, J=4.8, H-2'), 1.98 (1H, m, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.55 (1H, m, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 156.01 (C-6), 152.48 (C-2), 149.69 (C-4), 141.11 (C-8), 118.68 (C-5), 70.71 (C-2', d, J=13.8), 68.30 (CH<sub>2</sub>OP, d, J=6.9), 64.55 (C-4', d, J=158), 42.52 (C-1'), 25.85 (CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>, d, J=9.0). IR (KBr) 3351, 3169, 1660, 1601, 1256, 1063. MS (FAB) 314 (M+H, 100).

Anal. Calcd for: C <sub>11</sub> H <sub>16</sub> N <sub>5</sub> O <sub>4</sub> P · 1.5 H <sub>2</sub> O	C, 38.85;	H, 5.63;	N, 20.60.
Found:	C, 38.63;	H, 5.46;	N, 20.49.

## EXAMPLE 18

PMEA, bis-diethylamide

Mp 93-96°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.11 (1H, s, H-8), 8.07 (1H, s, H-2), 7.18 (2H, s, NH<sub>2</sub>), 4.31 (2H, t, J=4.8, H-1'), 3.85 (2H, t, J=4.8, H-2'), 3.68 (2H, d, J=8.1, H-4'), 2.70 (8H, m, CH<sub>3</sub>CH<sub>2</sub>), 0.86 (12H, t, J=7.0, CH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 155.98 (C-6), 152.33 (C-2), 149.63 (C-4), 141.04 (C-8), 118.75 (C-5), 70.30 (C-2', d, J=13.0), 66.30 (C-4', d, J=133), 42.63 (C-1'), 37.53 (CH<sub>3</sub>CH<sub>2</sub>, d, J=4.1), 13.93 (CH<sub>3</sub>, d, J=1.9). IR (KBr) 3370-2935, 2875, 1680, 1649, 1605, 1211. MS (FAB) 384 (M+H, 100).

Anal. Calcd for: C <sub>16</sub> H <sub>30</sub> N <sub>7</sub> O <sub>2</sub> P · 0.5 H <sub>2</sub> O	C, 48.96;	H, 7.96;	N, 24.99.
Found:	C, 48.85;	H, 7.77;	N, 24.92.

## EXAMPLE 19

PMEA, mono-(isopropyl ester) (sodium salt)

5 Mp 77-85°C turned to glass and melted over next 40°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.19 (1H, s, H-8), 8.13 (1H, s, H-2), 7.22 (2H, s, NH<sub>2</sub>), 4.30 (2H, t, J=4.4, H-1'), 4.10 (1H, m, OCH), 3.76 (2H, t, J=4.4, H-2'), 3.31 (2H, d, J=8.6, H-4'), 0.90 (6H, d, J=6.0, CH<sub>3</sub>). <sup>13</sup>C (d<sub>6</sub>-DMSO; 90 MHz) 155.90 (C-6), 152.35 (C-2), 149.54 (C-4), 141.39 (C-8), 118.53 (C-5), 70.23 (OCH, d, J=10), 68.70 (C-4', d, J=192), 65.55 (C-2', d, J=5), 42.72 (C-1'), 24.43 (CH<sub>3</sub>). IR (Film) 3321, 3163, 1647, 1601, 1578. MS (FAB) 338 (M+H, 70).

Anal. Calcd for: C <sub>11</sub> H <sub>17</sub> N <sub>5</sub> O <sub>4</sub> P <sub>1</sub> Na <sub>1</sub> · H <sub>2</sub> O	C, 37.18;	H, 5.38;	N, 19.71.
Found:	C, 37.11;	H, 5.49;	N, 19.71.

## EXAMPLE 20

PMEA, cyclic (2,2-dimethyl)propanyl diester

20 Mp 224-226°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.11 (2H, s, H-8, H-2), 7.21 (2H, s, NH<sub>2</sub>), 4.34 (2H, t, J=5.0, H-1'), 3.99 (2H, d, J=8.7, H-4'), 3.91 (2H, t, J=5.0, H-2'), 3.95-3.75 (4H, m, CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>), 1.06 (3H, s, CH<sub>3</sub>), 0.67 (3H, s, CH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO; 50 MHz) 155.89 (C-6), 152.33 (C-2), 149.53 (C-4), 140.86 (C-8), 118.57 (C-5), 76.67 (CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>CH<sub>2</sub>, d, J=6.8), 70.44 (C-2', d, J=13.7), 64.43 (C-4', d, J=157), 42.43 (C-1'), 31.70 (C(CH<sub>3</sub>)<sub>2</sub>, d, J=7.6), 21.05 (CH<sub>3</sub>), 19.46 (CH<sub>3</sub>). IR (KBr) 3417, 3324, 3152, 2970, 1668, 1650, 1602. MS (FAB) 342 (M+H, 100).

Anal. Calcd for: C <sub>13</sub> H <sub>20</sub> N <sub>5</sub> O <sub>4</sub> P · 0.25 H <sub>2</sub> O	C, 45.18;	H, 5.97;	N, 20.27.
Found:	C, 45.58;	H, 6.05;	N, 20.05.

## EXAMPLE 21

PMEA, mono-(3-hydroxypropanyl ester) (sodium salt)

35 <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.17 (1H, s, H-8), 8.11 (1H, s, H-2), 7.20 (2H, s, NH<sub>2</sub>), 5.11 (1H, t, OH), 4.28 (2H, t, J=4.7, H-1'), 3.76 (2H, t, J=4.7, H-2'), 3.64 (2H, q, J=6.6, CH<sub>2</sub>CH<sub>2</sub>OP), 3.41 (2H, d, J=8.0, H-4'), 3.35 (2H, t, J=6.2, HOCH<sub>2</sub>), 1.45 (2H, m, HOCH<sub>2</sub>CH<sub>2</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO; 50 MHz) 155.82 (C-6), 152.25 (C-2), 149.43 (C-4), 141.38 (C-8), 118.43 (C-5), 69.77 (C-2', d, J=10), 67.42 (C-4', d, J=152), 59.33 (CH<sub>2</sub>CH<sub>2</sub>OP, d, J=6), 56.88 (HOCH<sub>2</sub>), 42.60 (C-1'), 33.91 (HOCH<sub>2</sub>CH<sub>2</sub>, d, J=4). IR (KBr) 3412, 2956, 1647, 1604, 1482, 1421. MS (FAB) 354 (M+H, 17).

Anal. Calcd for: C <sub>11</sub> H <sub>17</sub> N <sub>5</sub> O <sub>5</sub> P <sub>1</sub> Na <sub>1</sub> · 2.5 H <sub>2</sub> O	C, 33.17;	H, 5.56;	N, 17.59.
Found:	C, 33.32;	H, 5.28;	N, 17.63.

## EXAMPLE 22

PMEA, mono-(octyl ester)

50 <sup>1</sup>H NMR (d<sub>5</sub>-pyridine) 9.47, 9.34 (2H, 2s, H-2, H-8), 5.46 (2H, t, J=4.5), 5.3-5.1 (6H, m, H-2', H-4', CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O), 2.68 (2H, m, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O), 2.33 (2H, m, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>O), 2.1 (8H, m, CH<sub>3</sub>(CH<sub>2</sub>)<sub>4</sub>), 1.79 (3H, t, J=6.5, CH<sub>3</sub>). IR (KBr) 3416, 2928, 1690, 1065. MS (FAB) 386 (M+H, 100).

Anal. Calcd for: $C_{16}H_{28}N_5O_4P \cdot H_2O \cdot Na \cdot 0.6 NaCl$	C, 41.59;	H, 6.54;	N, 15.15.
Found:	C, 41.80;	H, 6.87;	N, 15.02.

## EXAMPLE 23

10 PMEA, mono-(3-amino-2,2-dimethylpropyl amide)

$^1H$  NMR ( $D_2O$ ) 8.13 (1H, s, H-8), 8.11 (1H, s, H-2), 4.36 (2H, t, J=5, H-1'), 3.90 (2H, t, J=5, H-2'), 3.53 (2H, d, J=8.5, H-4'), 2.71 (2H, s,  $NH_2CH_2$ ) 2.07 (2H, d, J=9.4,  $CH_2NH$ ), 0.70 (6H, s,  $CH_3$ ).  $^{13}C$  NMR ( $D_2O$ ) 157.25 (C-6), 154.19 (C-2), 150.78 (C-4), 144.73 (C-8), 120.03 (C-5), 72.24 (C-2', d, J=12.5), 69.63 (C-4', d, J=143), 50.05 ( $CH_2NH$ ), 48.41 ( $H_2NCH_2$ ), 45.53 (C-1), 35.36 ( $C(CH_3)_2$ , d, J=4), 24.09 ( $CH_3$ ). IR (KBr) 3786, 3381, 1648, 1605, 1478. MS (FAB) 380 (M+H, 20). HR-MS (M+H)

Anal. Calcd for $C_{13}H_{23}N_7O_3P_1Na_1$ :	380.1576.
Found:	380.1567.

## 25 EXAMPLE 24

PMEA, mono-(hydroxy-2,2-dimethylpropyl ester)

$^1H$  NMR ( $d_6$ -DMSO) 8.14 (1H, s, H-8), 8.09 (1H, s, H-2), 7.16 (2H, s,  $NH_2$ ), 5.84 (1H, t, OH), 4.27 (2H, t, J=4.9, H-1'), 3.77 (2H, t, J=4.9, H-2'), 3.33 (2H, d, J=8.7, H-4'), 3.24 (2H, d, J=10,  $C(CH_3)_2CH_2OP$ ), 3.00 (2H, d,  $HOCH_2$ ), 0.63 (6H, s,  $CH_3$ ).  $^{13}C$  NMR ( $d_6$ -DMSO, 50 MHz) 155.84 (C-6), 152.21 (C-2), 149.45 (C-4), 141.26 (C-8), 118.48 (C-5), 69.71 (C-2', d, J=9.2), 68.27 ( $C(CH_3)_2CH_2OP$ , d, J=6.2), 67.48 (C-4', d, J=152), 65.93 ( $HOCH_2$ ), 42.57 (C-1'), 36.71 ( $C(CH_3)_2$ , d, J=2.5), 21.35 ( $CH_3$ ). IR (KBr) 3426, 2960, 2883, 1645, 1478, 1417. MS (FAB) 360 (M+H, 100).

Anal. Calcd. for $C_{13}H_{22}N_5O_5P \cdot 1.3 H_2O$ :	C, 40.77;	H, 6.48;	N, 18.29.
Found:	C, 40.96;	H, 6.16;	N, 17.95

## EXAMPLE 25

PMEA, cyclic (2,2-dimethyl-propanyl diamide)

$^1H$  NMR ( $d_6$ -DMSO) 8.11 (1H, s, H-8), 8.10 (1H, s, H-2), 7.18 (2H, s,  $NH_2$ ), 4.30 (2H, t, J=5.0, H-1'), 3.83 (2H, t, J=5.0, H-2'), 3.63 (2H, d, J=7.5, H-4'), 4.27 (2H, s, NH, NH), 2.65-2.40 (4H, m,  $CH_2C(CH_3)_2CH_2$ ), 0.98 (3H, s,  $CH_3$ ), 0.64 (3H, s,  $CH_3$ ).  $^{13}C$  NMR ( $d_6$ -DMSO) 156.01 (C-6), 152.42 (C-2), 149.60 (C-4), 141.24 (C-8), 118.68 (C-5), 70.35 (C-2', d, J=11.2), 68.53 (C-4', d, J=131), 52.72 ( $CH_2C(CH_3)_2CH_2$ , d, J=2.3), 42.78 (C-1'), 30.54 ( $C(CH_3)_2$ , d, J=5.6), 24.82 ( $CH_3$ ), 23.25 ( $CH_3$ ). IR (KBr) 3100, 2980, 2940, 1650, 1605. MS (FAB) 340 (M+H, 100). HR-MS (M+H)

Anal. Calcd for $C_{13}H_{22}N_7O_2P$ :	340.1651.
Found:	340.1647.

## EXAMPLE 26

PMEA, N,N'-dimethyl-cyclic propanyl diamide

<sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.08 (2H, s, H-8, H-2), 7.14 (2H, s, NH<sub>2</sub>), 4.28 (2H, br s, H-1'), 3.80 (2H, br s, H-2'), 3.73 (2H, dd, J=7.6, 2.8 H-4'), 2.85-2.60 (4H, m, CH<sub>3</sub>NCH<sub>2</sub>), 1.8-1.3 (2H, m, CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>), 2.36 (3H, d, J=3, NCH<sub>3</sub>), 2.33 (3H, d, J=3, NCH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 156.02 (C-6), 152.44 (C-2), 149.77 (C-4), 141.09 (C-8), 118.74 (C-5), 70.44 (C-2', d, J=14), 65.42 (C-4', d, J=164), 50.22 (NCH<sub>3</sub>), 42.85 (C-1'), 34.28 (CH<sub>3</sub>NCH<sub>2</sub>), 24.79 (CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>). IR (KBr) 3300, 3180, 2930, 2877, 1651, 1600. MS (methane/DCI) 340 (M+H, 100).

Anal. Calcd for C <sub>13</sub> H <sub>22</sub> N <sub>7</sub> O <sub>2</sub> P · 0.9 HCl:	C, 41.93;	H, 6.22;	N, 26.33.
Found:	C, 42.33;	H, 6.19;	N, 25.93.

HR-MS (M+H) Calcd for C <sub>13</sub> H <sub>22</sub> N <sub>7</sub> O <sub>2</sub> P:	340.1651.
Found:	340.1649.

## EXAMPLE 27

PMEA, mono-(N,N'-diethylacetamide ester)

Mp 189-191°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.16 (1H, s, H-8), 8.14 (1H, s, H-2), 7.55 (2H, s, NH<sub>2</sub>), 4.80 (2H, d, J=9.0, C(O)CH<sub>2</sub>O), 4.31 (2H, t, J=5.0, H-1'), 4.03 (2H, t, J=5.0, H-2'), 3.74 (2H, d, J=8.5, H-4'), 3.22 (2H, q, J=7, CH<sub>3</sub>CH<sub>2</sub>), 3.16 (2H, q, J=7, CH<sub>3</sub>CH<sub>2</sub>), 1.01 (3H, t, J=7, CH<sub>3</sub>), 1.01 (3H, t, J=7, CH<sub>3</sub>). <sup>13</sup>C NMR (CF<sub>3</sub>CO<sub>2</sub>D; 90 MHz) 166.10 (C=O), 150.04, 148.67 (C-6, C-4), 144.74, 144.55 (C-2, C-8), 117.96 (C-5), 70.05 (C-2', d, J=10), 65.37 (C-4', d, J=162), 62.87 (C(O)CH<sub>2</sub>, d, J=5), 43.44 (C-1'), 14.06 (CH<sub>3</sub>), 12.91 (CH<sub>3</sub>). IR (KBr) 3392, 3093, 1692, 1650, 1515. MS (methane/DCI) 500 (M+H, 30), 132 (100). HR-MS (M+H)

Anal. Calcd for C <sub>14</sub> H <sub>23</sub> N <sub>5</sub> O <sub>5</sub> P:	387.1546.
Found:	387.1543.

## EXAMPLE 28

PMEA, mono-(acetic acid ester)

Mp 197-200°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.19 (1H, s, H-8), 8.17 (1H, s, H-2), 7.75 (2H, s, NH<sub>2</sub>), 4.34 (2H, d, J=4, C(O)CH<sub>2</sub>O), 4.32 (2H, t, J=5, H-1'), 3.86 (2H, t, J=5, H-2'), 3.71 (2H, d, J=8, H-4'). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 177.19 (C=O, d, J=7), 156.84 (C-6), 153.72 (C-2), 150.03 (C-4), 144.05 (C-8), 119.44 (C-5), 71.66 (C-2', d, J=11), 67.39 (C-4', d, J=157), 64.90 (C(O)CH<sub>2</sub>O, d, J=6), 44.59 (C-1'). IR (KBr) 3366, 3109, 1690, 1611, 1516, 1415. MS (FAB) 332 (M+H, 55).

Anal. Calcd for C <sub>10</sub> H <sub>14</sub> N <sub>5</sub> O <sub>6</sub> P · 0.3 H <sub>2</sub> O:	C, 35.74;	H, 4.38;	N, 20.85.
Found:	C, 35.41;	H, 4.43;	N, 20.60.

## EXAMPLE 29

PMEA, di-(butylacetate ester)

5 Mp 78-80°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.11 (1H, s, H-8), 8.06 (1H, s, H-2), 7.18 (2H, s, NH<sub>2</sub>), 4.62 (4H, d, J=11, C(O)CH<sub>2</sub>OP), 4.31 (2H, t, J=5.0, H-1'), 4.07 (4H, t, J=7, CH<sub>2</sub>OC(O)), 4.00 (2H, d, J=8, H-4'), 3.90 (2H, t, J=5, H-2'), 1.54 (4H, apparent quintet, J=7, CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>), 1.31 (4H, apparent hexet, J=7.7, CH<sub>3</sub>CH<sub>2</sub>), 0.86 (6H, t, J=7, CH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 168.16 (C=O, d, J=4.7), 156.03 (C-6), 152.44 (C-2), 149.59 (C-4), 141.10 (C-8), 118.65 (C-5), 70.58 (C-2', d, J=10), 64.70 (CH<sub>2</sub>OC(O)), 64.19 (C-4', d, J=165), 62.05 (CH<sub>2</sub>OP, d, J=6), 42.45 (C-1'), 30.10 (CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>), 18.53 (CH<sub>3</sub>CH<sub>2</sub>), 13.56 (CH<sub>3</sub>). IR (KBr) 3339, 3158, 2994, 2962, 1764, 1662, 1600. MS (methane/DCI) 502 (M+H, 100).

15

Anal. Calcd for C <sub>20</sub> H <sub>32</sub> N <sub>5</sub> O <sub>8</sub> P:	C, 47.90;	H, 6.43;	N, 13.97.
Found:	C, 47.94;	H, 6.40;	N, 13.90.

## EXAMPLE 30

PMEA, di-(ethylacetate ester)

20 Mp 82-84°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.11 (1H, s, H-8), 8.06 (1H, s, H-2), 7.16 (2H, s, NH<sub>2</sub>), 4.59 (4H, d, J=11, C(O)CH<sub>2</sub>O), 4.30 (2H, t, J=5.0, H-1'), 4.13 (4H, q, J=7.0, CH<sub>3</sub>CH<sub>2</sub>), 4.00 (2H, d, J=8.0, H-4'), 3.98 (2H, t, J=5.0, H-2'), 1.18 (6H, t, J=7.0, CH<sub>3</sub>). <sup>13</sup>C NMR (D<sub>2</sub>O) 171.44 (C=O, d, J=5), 156.90 (C-6), 153.85 (C-2), 150.56 (C-4), 144.66 (C-8), 119.86 (C-5), 73.02 (C-2', d, J=10.5), 66.12 (C-4', d, J=166), 64.85 (CH<sub>3</sub>CH<sub>2</sub>), 64.75 (C(O)CH<sub>2</sub>O), 45.57 (C-1') 15.22 (CH<sub>3</sub>). IR (KBr) 3296, 3122, 1764, 1667, 1602. MS (methane/DCI) 446 (M+H, 100).

30

Anal. Calcd for C <sub>16</sub> H <sub>24</sub> N <sub>5</sub> O <sub>8</sub> P:	C, 43.15;	H, 5.43;	N, 15.72.
Found:	C, 43.04;	H, 5.33;	N, 15.58.

## EXAMPLE 31

PMEA, mono-(phenyl ester) (sodium salt)

35 Mp 223-228°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.14 (1H, s, H-8), 8.13 (1H, s, H-2), 7.50 (2H, s, NH<sub>2</sub>), 7.25 (2H, t, J=8, ArH), 7.07 (1H, t, J=8, ArH), 7.01 (2H, d, J=8, ArH), 4.33 (2H, t, J=5, H-1'), 3.89 (2H, t, J=5, H-2'), 3.73 (2H, d, J=8, H-4'). <sup>13</sup>C NMR (D<sub>2</sub>O; Partial spectrum) 131.46, 126.06 (ArC), 122.27 (ArC, d, J=3.5), 72.27 (C-2; d, J=12), 67.68 (C-4', d, J=160), 46.08 (C-1'). IR (KBr) 3389, 3068, 1693, 1594. MS (FAB) 350 (M+H, 40).

45

Anal. Calcd for C <sub>14</sub> H <sub>16</sub> N <sub>5</sub> O <sub>4</sub> P · H <sub>2</sub> O · 0.45 Na:	C, 44.45;	H, 4.81;	N, 18.51.
Found:	C, 44.45;	H, 4.45;	N, 18.45.

50

## EXAMPLE 32

PMEA, di-(phenyl ester)

55 Mp 103-114°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.15 (1H, s, H-8), 8.11 (1H, s, H-2), 7.40 (2H, s, NH<sub>2</sub>), 7.34 (4H, t, J=7, ArH), 7.20 (2H, t, J=7, ArH), 7.04 (4H, t, J=7, ArH), 4.38 (2H, t, J=5, H-1'), 4.24 (2H, d, J=8, H-4'), 3.98 (2H, t, J=5, H-2'). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 155.51 (C-6), 151.77 (C-2), 149.57 (C-4), 141.46 (C-8), 130.02, 125.49, (ArC), 120.56 (ArC, d, J=4), 118.71 (C-5), 70.58 (C-2', d, J=12), 63.52 (C-4', d, J=164), 42.68 (C-1'). IR (KBr) 3270, 3100, 1675, 1646,

1601, 1490. MS (FAB) 426 (M+H, 100).

Anal. Calcd for $C_{20}H_{20}N_5O_4P \cdot 0.25 H_2O$ :	C, 55.87;	H, 4.81;	N, 16.29.
Found:	C, 55.80;	H, 4.65;	N, 15.98.

## EXAMPLE 33

PMEA, mono-(N,N-diisopropylacetamide ester) (sodium salt)

Mp 219-221°C.  $^1H$  NMR ( $d_6$ -DMSO) 8.14 (1H, s, H-8), 8.13 (1H, s, H-2), 7.37 (2H, s,  $NH_2$ ), 4.45 (2H, d, J=9,  $CH_2OP$ ), 4.31 (2H, t, J=5, H-1'), 3.88 (2H, t, J=5, H-2'), 3.74 (2H, d, J=8, H-4'), 3.43 (2H, m,  $CH(CH_3)_2$ ), 1.26 (6H, d, J=6,  $CH_3$ ), 1.08 (6H, d, J=6,  $CH_3$ ).  $^{13}C$  NMR ( $d_6$ -DMSO/ $D_2O$ ) 170 (C=O), 156.90 (C-6), 153.89 (C-2), 150.35 (C-4), 144.29 (C-8), 119.68 (C-5), 71.89 (C-2', d, J=12), 67.81 (C-4', d, J=158), 65.25 ( $CH_2OP$ , d, J=5), 49.72 ( $CH(CH_3)_2$ ), 47.30 ( $CH(CH_3)_2$ ), 45.00 (C-1'), 21.21 ( $CH_3$ ). IR (KBr) 3425, 2969, 1691, 1643, 1515. MS (FAB) 415 (M+H, 100).

Anal. Calcd for $C_{16}H_{27}N_6O_5P \cdot 0.67 H_2O \cdot 0.5 Na$ :	C, 43.87;	H, 6.52;	N, 19.19.
Found:	C, 43.92;	H, 6.17;	N, 18.79.

## EXAMPLE 34

PMEA, di-(p-nitrobenzyl ester)

Mp 190-193°C.  $^1H$  NMR ( $d_6$ -DMSO) 8.16 (4H, d, J=8, ArH), 8.09 (1H, s, H-8), 8.08 (1H, s, H-2), 7.51 (4H, d, J=8, ArH), 7.17 (2H, s,  $NH_2$ ), 5.10 (4H, d, J=8,  $ArCH_2O$ ), 4.32 (2H, t, J=5, H-1'), 4.07 (2H, d, J=8, H-4'), 3.90 (2H, t, J=5, H-2'),  $^{13}C$  NMR ( $d_6$ -DMSO) 155.97 (C-6), 152.94 (C-2), 149.62 (C-4), 147.19, 143.96 (ArC), 141.13 (C-8), 128.15, 123.56 (ArC), 118.65 (C-5), 70.62 (C-2', d, J=7), 65.86 ( $ArCH_2O$ , d, J=6), 63.75 (C-4', d, J=162), 42.49 (C-1'). IR (KBr) 3420, 3268, 3110, 1674, 1642, 1604. MS (FAB) 544 (M+H, 60).

Anal. Calcd for $C_{22}H_{22}N_7O_8P$ :	C, 48.63;	H, 4.09;	N, 18.05.
Found:	C, 48.61;	H, 4.01;	N, 18.04.

## EXAMPLE 35

PMEA, mono-(p-nitrobenzyl ester) (sodium salt)

Mp 230-240°C.  $^1H$  NMR ( $d_6$ -DMSO) 8.19 (2H, d, J=8.6, ArH), 8.12 (1H, s, H-8), 8.11 (1H, s, H-2), 7.54 (2H, d, J=8.6, ArH), 4.93 (2H, d, J=7.7,  $ArCH_2O$ ), 4.63 (2H, t, J=5, H-1'), 4.31 (2H, t, J=5, H-2'), 3.72 (2H, d, J=8.6, H-4'). IR (KBr) 3742, 1930, 1692, 1606, 1518. MS (FAB) 409 (M+H, 27).

Anal. Calcd for $C_{15}H_{17}N_6O_6P \cdot 0.75 H_2O \cdot 0.5 Na$ :	C, 41.58;	H, 4.30;	N, 19.40.
Found:	C, 41.37;	H, 3.92;	N, 19.03.

## EXAMPLE 36

PMEA, di-(2,2,2-trichloroethyl ester)

5 Mp 155-157°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.11 (1H, s, H-8), 8.08 (1H, s, H-2), 7.16 (2H, s, NH<sub>2</sub>), 4.68 (2H, d, J=7, CCl<sub>3</sub>CH<sub>2</sub>), 4.67 (2H, d, J=7, CCl<sub>3</sub>CH<sub>2</sub>), 4.34 (2H, t, J=5, H-1'), 4.18 (2H, d, J=8, H-4'), 3.95 (2H, t, J=5, H-2'). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 156.09 (C-6), 152.59 (C-2), 149.71 (C-4), 141.28 (C-8), 118.75 (C-5), 95.42 (CCl<sub>3</sub>, d, J=8.6), 75.48 (CCl<sub>3</sub>CH<sub>2</sub>, d, J=5.7), 70.92 (C-2', d, J=7), 63.99 (C-4', d, J=163), 42.72 (C-1'). IR (KBr) 3372, 3334, 3210, 1658, 1604, 1576. MS (methane/DCI) 536 (100), 534 (50), 192 (95).

Anal. Calcd for C <sub>12</sub> H <sub>14</sub> N <sub>5</sub> O <sub>4</sub> PCl <sub>6</sub> :	C, 26.89;	H, 2.63;	N, 13.07.
Found:	C, 26.85;	H, 2.55;	N, 12.86.

## EXAMPLE 37

PMEA, mono-(2,2,2-trichloroethyl ester)

20 Mp 218-225°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.51 (2H, s, NH<sub>2</sub>), 8.30, 8.24 (2H, 2s, H-8, H-2), 4.36 (2H, t, J=5, H-1'), 4.33 (2H, d, J=6, Cl<sub>3</sub>CCH<sub>2</sub>), 3.72 (2H, d, J=8, C-4'), 3.91 (2H, t, J=5, H-2'). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 153.03 (C-6), 148.91 (C-2), 148.22 (C-4), 142.78 (C-8), 118.27 (C-5), 97.05 (CCl<sub>3</sub>), 75.67 (CCl<sub>3</sub>CH<sub>2</sub>, d, J=5), 69.99 (C-2', d, J=10), 66.17 (C-4', d, J=159), 43.12 (C-1'). IR (KBr) 3424, 1930, 1690, 1614, 1514, 1414. MS (methane/DCI) 404 (M+H, 1), 136 (40), 113 (100).

Anal. Calcd for C <sub>10</sub> H <sub>13</sub> N <sub>5</sub> O <sub>4</sub> PCl <sub>3</sub> · 0.3 CCl <sub>3</sub> CH <sub>2</sub> OH:	C, 28.34;	H, 3.05;	N, 15.59.
Found:	C, 28.19;	H, 3.17;	N, 15.59.

## EXAMPLE 38

PMEA, di-(benzoyloxymethyl ester)

35 Mp 49-52°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.09 (1H, s, H-8), 7.99 (1H, s, H-2), 7.92 (4H, d, J=7, ArH), 7.67 (2H, t, J=7.5, ArH), 7.49 (2H, t, J=7.5, ArH), 7.18 (2H, s, NH<sub>2</sub>), 5.82 (4H, d, J=13, OCH<sub>2</sub>O), 4.22 (2H, t, J=5, H-1'), 4.04 (2H, d, J=8, H-4'), 3.82 (2H, d, J=5, H-2'). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 164.35 (C=O), 156.02 (C-6), 152.45 (C-2), 149.55 (C-4), 140.99 (C-8), 134.22 (ArH), 129.60 (ArH), 128.98 (ArH), 128.35 (ArH), 118.70 (C-5), 70.62 (C-2', d, J=11.5), 64.17 (C-4', d, J=163), 42.29 (C-1'). IR (KBr) 3328, 3182, 1739, 1644, 1602. MS (FAB) 542 (M+H, 45).

Anal. Calcd for C <sub>24</sub> H <sub>24</sub> N <sub>5</sub> O <sub>6</sub> P · 0.66 H <sub>2</sub> O:	C, 52.09;	H, 4.61;	N, 12.65.
Found:	C, 52.09;	H, 4.36;	N, 12.37.

## EXAMPLE 39

PMEA, di-(p-trifluoromethyl benzyl ester)

50 Mp 115-125°C. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) 8.18 (1H, s, H-8), 8.17 (1H, s, H-2), 7.66 (4H, d, J=8, ArH), 7.47 (4H, d, J=8, ArH), 7.57 (2H, s, NH<sub>2</sub>), 5.09 (4H, d, J=8, ArCH<sub>2</sub>), 4.35 (2H, t, J=5, H-1'), 4.04 (2H, d, J=8, H-4'), 3.91 (2H, t, J=5, H-2'). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 154.99 (C-6), 151.13 (C-2), 149.44 (C-4), 141.7 (C-8), 141.12 (ArC), 128.63 (CF<sub>3</sub>-ArC, q, J=31.8), 127.93, 125.31 (ArC), 124.17 (CF<sub>3</sub>, q, J=275), 118.53 (C-5), 70.46 (C-2', d, J=11), 66.14 (ArCH<sub>2</sub>, d, J=5.5), 63.78 (C-4', d, J=161), 42.61 (C-1'). IR (KBr) 3292, 3118, 1670, 1602, 1476. MS (FAB) 590 (M+H, 100).



Anal. Calcd for $C_{24}H_{22}N_5O_4PF_6 \cdot 0.5 H_2O$ :	C, 48.17;	H, 3.87;	N, 11.70.
Found:	C, 47.81;	H, 3.55;	N, 11.30.

## EXAMPLE 40

10 PMEA, mono-(2,2-difluoro-3-hydroxypropyl ester)

$^1H$  NMR ( $d_6$ -DMSO) 8.20 (2H, s, H-8, H-2), 7.80 (2H, s,  $NH_2$ ), 4.34 (2H, t,  $J=5.0$ , H-1'), 4.04 (2H, dt,  $J=13.2$ , 7.9),  $CF_2CH_2OP$ ), 3.87 (2H, t,  $J=5.0$ , H-2'), 3.70 (2H, d,  $J=8.0$ , H-4'), 3.60 (2H, t,  $J=13$ ,  $HOCH_2$ ).  $^{13}C$  NMR ( $D_2O/NaOD$ ) 157.34 (C-6), 154.24 (C-2), 150.67 (C-4), 144.72 (C-8), 123.54 ( $CF_2$ , t,  $J=30$ ), 120.12 (C-5), 72.40 (C-2', d,  $J=12$ ), 67.75 (C-4', d,  $J=159$ ), 64.94 ( $CF_2CH_2OP$ , dt,  $J=30$ , 5), 63.28 ( $HOCH_2$ , d,  $J=27$ ), 45.49 (C-1'). IR (KBr) 3310, 3112, 1694, 1602, 1514. MS (FAB) 368 (M+H, 55). HR-MS (M+H).

Anal. Calcd for $C_{11}H_{16}N_5O_5F_2P$ :	368.0935.
Found:	368.0930.

## EXAMPLE 41

25 PMEA, mono-(p-trifluoromethylbenzyl ester)

$^1H$  NMR ( $d_6$ -DMSO) 8.13 (2H, s, H-8, H-2), 7.69 (2H, d,  $J=8$ , ArH), 7.49 (2H, d,  $J=8$ , ArH), 7.34 (2H, s,  $NH_2$ ), 4.92 (2H, d,  $J=8$ ,  $ArCH_2O$ ), 4.32 (2H, t,  $J=5$ , C-1'), 3.87 (2H, t,  $J=5$ , H-2'), 3.75 (2H, d,  $J=8$ , H-4'). IR (KBr) 3062, 1696, 1602, 1514, 1418. MS (FAB) 432 (M+H, 80). HR-MS (M+H).

Anal. Calcd for $C_{16}H_{17}N_5O_4PF_3$ :	432.1048.
Found:	432.1039.

## EXAMPLE 42

40 PMEA, dibutylamide

Mp 117-119°C.  $^1H$  NMR ( $d_6$ -DMSO) 8.12 (2H, s, H-8, H-2), 7.19 (2H, s,  $NH_2$ ), 4.29 (2H, t,  $J=5$ , H-1'), 3.82 (2H, t,  $J=5$ , H-2'), 3.83 (2H, s, NH), 3.52 (2H, d,  $J=8$ , H-4'), 2.64 (4H, m,  $CH_2NH$ ), 1.24 (8H, m,  $CH_3CH_2CH_2$ ), 0.80 (6H, t,  $J=7$ ,  $CH_3$ ).  $^{13}C$  NMR ( $d_6$ -DMSO) 155.98 (C-6), 152.61 (C-2), 149.71 (C-4), 141.52 (C-8), 118.65 (C-5), 70.46 (C-2', d,  $J=11$ ), 67.28 (C-4', d,  $J=131$ ), 42.83 (C-1'), 39.22 ( $NHCH_2$ ), 34.10 ( $NHCH_2CH_2$ ), 19.59 ( $CH_3CH_2$ ), 13.92 ( $CH_3$ ). IR 3278, 3242, 2952, 2928, 2872, 1682, 1608. MS (FAB) 384 (M+H, 100).

Anal. Calcd for $C_{16}H_{30}N_7O_2P$ :	C, 50.12;	H, 7.89;	N, 25.57.
Found:	C, 49.77;	H, 7.79;	N, 25.30.

## EXAMPLE 43

55 PMEA, di-(2-methylpropyl ester)

Mp 109-110°C.  $^1H$  NMR ( $d_6$ -DMSO) 8.10 (1H, s, H-8), 8.05 (1H, s, H-2), 7.19 (2H, s,  $NH_2$ ), 4.31 (2H, t,  $J=5.0$ , H-1'),

3.87(2H,t,J=5.0,H-2'), 3.85(2H,d,J=8.5,H-4'), 3.61(4H,dt,J=6.8,1.4,CH<sub>2</sub>OP), 1.72(2H, apparent heptet, J=6.7,CH), 0.77(12H,d,J=6.7,CH<sub>3</sub>). <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) 156.04(C-6), 152.42(C-2), 149.60(C-4), 141.05(C-8), 118.69(C-5), 71.42(CH<sub>2</sub>OP, d, J=6.7), 70.36(C-2',d,J=11.6), 63.65(C-4',d,J=163), 42.52(C-1'), 28.72(CH,d,J=5.7), 18.45(CH<sub>3</sub>). IR(KBr) 3286, 3104, 2960, 1670, 1600. MS(FAB) 386(M+H, 100).

Anal. Calcd for C <sub>16</sub> H <sub>28</sub> N <sub>5</sub> O <sub>4</sub> P <sub>1</sub> :	C, 49.86;	H, 7.32;	N, 18.17.
Found:	C, 49.81;	H, 7.26;	N, 18.11

## EXAMPLE 44

## PMEA, di-(3-methylbutyl ester)


Mp 94-98°C. <sup>1</sup>H NMR(CDCl<sub>3</sub>) 8.30(1H, S, H-8) 7.94(1H, S, H-2), 6.21(2H, S, NH<sub>2</sub>), 4.37(2H, t, J=5.0, H-1'), 4.01(4H,dt, J=6.8, 6.8, CH<sub>2</sub>OP), 3.91(2H, t, J=5.0, H-2'), 3.75(2H,d,J=8.0,H-4'), 1.63(2H, apparent heptet, J=6.6, CH), 1.47(4H,dt, J=6.7, 6.7, CH<sub>2</sub>CH<sub>2</sub>OP), 0.84(12H,d,J=6.5,CH<sub>3</sub>). <sup>13</sup>C NMR (CDCl<sub>3</sub>) 155.28(C-6), 152.38(C-2), 150.38(C-4), 141.70(C-8), 119.76(C-5), 71.13(C-2',d,J=10.0), 65.17(C-4',d,J=166), 65.02 (CH<sub>2</sub>OP,d,J=6.8), 43.46(C-1'), 39.19 (CH<sub>2</sub>CH<sub>2</sub>OP,d,J=5.7), 24.50(CH), 22.31(CH<sub>3</sub>), 22.29(CH<sub>3</sub>). IR(KBr) 3282, 3106, 2958, 1672, 1600, 1478. MS(methane/DCI) 414(M+H,100).

Anal. Calcd. for C <sub>18</sub> H <sub>32</sub> N <sub>5</sub> O <sub>4</sub> P <sub>1</sub> · 0.75H <sub>2</sub> O:	C, 50.63;	H, 7.91;	N, 16.40.
Found:	C, 50.67;	H, 7.66;	N, 16.26.

## Claims

Claims for the following Contracting States : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE


1. A compound having the structural formula I



FORMULA I

wherein

B represents adenine (A), cytosine (C), guanine (G), thymine (T), uracil (U), 2,6-diamino purine (DAP), hypoxanthine (Hx),



wherein

Q is independently chosen from H, Cl,  $\text{NHR}^5$ ,  $\text{NR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH or  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  and  $\text{R}^2$  are identical or different and independently of one another are each  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  or  $\text{N(R}^5)_2$ ;  $\text{R}^1$  and  $\text{R}^2$  optionally being linked with each other to form a cyclic group, or  $\text{R}^1$  or  $\text{R}^2$  optionally being linked to  $\text{R}^3$  to form a cyclic group;

$\text{R}^3$  represents  $\text{C}_1$ - $\text{C}_{20}$  alkylene which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen; or  $\text{R}^3$  is  $\text{CH}(\text{CH}_2\text{OR}^6)\text{CH}_2$ , whereby  $\text{R}^1$  and  $\text{R}^2$  each independently may additionally represent OH, and  $\text{R}^6$  is a hydrolyzable ester group;


$\text{R}^4$  represents a physiologically hydrolyzable ester group selected from  $\text{CH}_2\text{C(O)NR}^5_2$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5)\text{OC(O)R}^5$  ( $\text{R}$ ; S; or RS stereochemistry),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$ , or  $\text{CH}_2\text{OR}^5$ ; or  $\text{R}^4$  represents  $\text{C}_4$ - $\text{C}_{20}$  alkyl, aryl-alkyl or aryl which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen provided that  $\text{R}^1$  and  $\text{R}^2$  are not simultaneously alkoxy;

$\text{R}^5$  represents  $\text{C}_1$ - $\text{C}_{20}$  alkyl, aryl or aryl-alkyl which may be substituted or unsubstituted by substituents independently selected from the group consisting of hydroxy and halogen;

pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;


provided that those compounds of the above formula I are excluded wherein

B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil,  $\text{R}^3$  is



wherein  $\text{alk}_1$  is bonded to B,  $\text{alk}_1$ ,  $\text{alk}_2$  and  $\text{alk}_3$  are independently selected from a chemical bond or  $\text{C}_1$ - $\text{C}_4$  alkylene,  $\text{R}^4$  is hydrogen or  $\text{C}_1$ - $\text{C}_4$  alkyl and  $\text{Q}^1$  is hydrogen or hydroxyl, and  $\text{R}^1$  and  $\text{R}^2$  are unsubstituted  $\text{C}_4$ - $\text{C}_6$  alkoxy, phenoxy or phenyl- $\text{C}_1$ - $\text{C}_4$  alkoxy.

## 2. The compound of claim 1 which has the general structural formula II



FORMULA II


wherein

B,  $\text{R}^1$  and  $\text{R}^2$  are as described in claim 1, provided that when Q is  $\text{NCHN(R}^5)_2$ , then  $\text{R}^5$  is not  $\text{CH}_3$ ;

X represents hydrogen,  $\text{CH}_2\text{OR}^6$  ( $\text{R}$ ; S; or RS stereochemistry), or substituted or unsubstituted lower alkyl, in

particular methyl or hydroxymethyl; when X is  $\text{CH}_2\text{OR}^6$ ,  $\text{R}^1$  and  $\text{R}^2$ , may additionally be independently chosen from OH; and  
 $\text{R}^6$  is a hydrolyzable ester group.

3. A compound having the general structural formula III



FORMULA III

wherein

B, and  $\text{R}^1$  are as previously described in claim 1;

X represents hydrogen,  $\text{CH}_2\text{OR}^6$  ( $\text{R}^6$  S; or R S stereochemistry) or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is  $\text{CH}_2\text{OR}^6$ ,  $\text{R}^1$  may additionally be OH; and  $\text{R}^6$  is a hydrolyzable ester group;

$\text{R}^7$  represents OH,  $\text{NH}_2$ ,  $\text{NHR}^5$ , or  $\text{NR}^5_2$ ; and

$\text{R}^5$  is as described in claim 1; pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;


provided that those compounds of the above formula III are excluded wherein

B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil,

X is  $\text{alk}_2\text{-Q}^1$  wherein  $\text{alk}_2$  is selected from a chemical bond or  $\text{C}_1\text{-C}_4$  alkylene,  $\text{Q}^1$  is hydrogen or hydroxyl, and

$\text{R}^1$  and  $\text{R}^7$  are OH, unsubstituted  $\text{C}_4\text{-C}_6$  alkoxy, phenoxy or phenyl- $\text{C}_1\text{-C}_4$  alkoxy.

4. The compound of claim 1 which has the general structural formula IV



FORMULA IV

wherein

$\text{R}^8$  and  $\text{R}^9$  are identical or different and independently of one another are each  $\text{NR}^{12}$ , or oxygen;

$\text{R}^{10}$  and  $\text{R}^{11}$  are identical or different and independently of one another are each hydrogen, or  $\text{R}^5$ ;


$\text{R}^{12}$  represents hydrogen or a lower alkyl;

m and n are identical or different and independently of one another are each 0 or 1;

B and  $\text{R}^5$  are as described in claim 1; and

X is as described in claim 2.

5. The compound which has the general structural formula V



\* stereochemistry is R; S; or RS

FORMULA V

wherein

$R^{13}$  represents  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$ , or OH, provided that  $R^{13}$  is not OH when B is A or C; and B,  $R^4$ , and  $R^5$  are as described in claim 1;

pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

6. The compound of claim 1, wherein  $R^4$  is  $C_4$ - $C_{20}$  alkyl, aryl or aryl-alkyl which may be unsubstituted or substituted by substituents independently selected from hydroxy and halogen.

7. The compound of claim 1 wherein  $R^1$  and  $R^2$  are  $OR^4$ , and  $R^4$  represents a physiologically hydrolyzable ester group selected from  $CH_2OC(O)R^5$ ; and  $CH(R^5)OC(O)R^5$  (R;S; or R,S stereochemistry); or aryl wherein aryl is substituted by substituents independently selected from hydroxy and halogen.

8. The compound of claim 1 which is

PMEA, di-(propionyloxymethyl ester);  
 PME, di-(isobutyryloxymethyl ester);  
 PME, bis-diethylamide;  
 PME, di-(butylacetate ester);  
 PME, di-(ethylacetate ester);  
 PME, di-(benzyloxymethyl ester);  
 PME, dibutylamide;  
 PME, di-(2-methylpropyl ester); or  
 PME, di-(3-methylbutyl ester).

9. The compound of claim 1 which is

PME, di-(pivaloyloxymethyl ester).

10. The compound of claim 3 which is

PME, mono-(pivaloyloxymethyl ester);  
 PME, mono-(octyl ester);  
 PME, mono-(hydroxy-2,2-dimethylpropyl ester);  
 PME, mono-(2,2,2-trichloroethyl ester); or  
 PME, mono-(2,2-difluoro-3-hydroxypropyl ester).

11. The compound of claim 4 which is

PME, cyclic propanyl diester;  
 PME, cyclic (2,2-dimethyl)propanyl diester;  
 PME, cyclic (2,2-dimethyl)propanyl diamide;  
 PME, N,N'-dimethyl-cyclic propanyl diamide.

12. The compound which is

PME, di-(isopropyl ester);

PMEHx, mono-(isopropyl ester);  
 6-chloro-9-(2-phosphonylmethoxy)ethylpurine,  
 di-(isopropyl ester);  
 9-(2-phosphonylmethoxy)ethylpurine, di-(isopropyl ester);  
 5   PMEA, mono-(isopropyl ester);  
 PMEA, (mono-isopropyl, mono-pivaloyloxymethyl) ester; or  
 PMEA, (mono-isopropyl, mono-phenyl) ester;  
 pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

10   13. The compound which is

PMEA, di-(phenyl ester);  
 PMEA, di-(p-nitrobenzyl ester);  
 PMEA, di-(2,2,2-trichloroethyl ester);  
 15   PMEA, di-(p-trifluoromethylbenzyl ester);  
 PMEA, mono-(choline ester);  
 PMEA, (mono-N,N-diethylacetamide, mono-pivaloyloxymethyl) ester;  
 PMEA, mono-(3-hydroxypropargyl ester);  
 PMEA, mono-(3-amino-2,2-dimethylpropyl amide);  
 20   PMEA, mono-(N,N-diethylacetamide ester);  
 PMEA, mono-(acetic acid ester);  
 PMEA, mono-(N,N-diisopropylacetamide ester);  
 PMEA, mono-(p-nitrobenzyl ester); or  
 PMEA, mono-(p-trifluoromethylbenzyl ester);  
 25   PMEA, mono-(phenyl ester);  
 pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

14. A process for producing the compound of claim 1, 2, 12, or 13 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or alkylation of the phosphonate with the appropriate alkyl halide; or hydrolysis of the diethers or diamines.

15. A process for producing the compound of claim 3, 12, or 13 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or hydrolysis of the diethers or diamines.

35   16. A process for producing the compound of claim 4 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol.

17. A process for producing the compound of claim 5 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or alkylation of the phosphonate with the appropriate alkyl halide.

18. The use of a compound as defined in anyone of claims 1 to 13 for preparing a pharmaceutical composition for the treatment of viral infection in a mammal.


45   19. The use of a compound as defined in anyone of claims 1 to 13 for preparing a pharmaceutical composition for inhibiting growth of a tumor in a mammal.

20. A pharmaceutical composition which comprises at least one compound of claims 1 to 13 in association with a pharmaceutically acceptable substantially nontoxic carrier or excipient.

50   Claims for the following Contracting State : ES

1. A process for producing a compound having the structural formula I


55



FORMULA I

wherein

B represents adenine (A), cytosine (C), guanine (G), thymine (T), uracil (U), 2,6-diamino purine (DAP), hypoxanthine (Hx),



wherein

Q is independently chosen from H, Cl,  $\text{NHR}^5$ ,  $\text{NR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH or  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  and  $\text{R}^2$  are identical or different and independently of one another are each  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  or  $\text{N(R}^5)_2$ ;  $\text{R}^1$  and  $\text{R}^2$  optionally being linked with each other to form a cyclic group, or  $\text{R}^1$  or  $\text{R}^2$  optionally being linked to  $\text{R}^3$  to form a cyclic group;

$\text{R}^3$  represents  $\text{C}_1$ - $\text{C}_{20}$  alkylene which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen; or  $\text{R}^3$  is  $\text{CH(CH}_2\text{OR}^6)\text{CH}_2$ , whereby  $\text{R}^1$  and  $\text{R}^2$  each independently may additionally represent OH, and  $\text{R}^6$  is a hydrolyzable ester group;


$\text{R}^4$  represents a physiologically hydrolyzable ester group selected from  $\text{CH}_2\text{C(O)NR}^5_2$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5\text{)OC(O)R}^5$  (R; S; or RS stereochemistry),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$ , or  $\text{CH}_2\text{OR}^5$ ; or  $\text{R}^4$  represents  $\text{C}_4$ - $\text{C}_{20}$  alkyl, aryl-alkyl or aryl which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen provided that  $\text{R}^1$  and  $\text{R}^2$  are not simultaneously alkoxy;

$\text{R}^5$  represents  $\text{C}_1$ - $\text{C}_{20}$  alkyl, aryl or aryl-alkyl which may be substituted or unsubstituted by substituents independently selected from the group consisting of hydroxy and halogen;

pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;

provided that those compounds of the above formula I are excluded wherein


B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil,  $\text{R}^3$  is



wherein  $\text{alk}_1$  is bonded to B,  $\text{alk}_1$ ,  $\text{alk}_2$  and  $\text{alk}_3$  are independently selected from a chemical bond or  $\text{C}_1$ - $\text{C}_4$  alkylene,  $\text{R}^a$  is hydrogen or  $\text{C}_1$ - $\text{C}_4$  alkyl and  $\text{Q}^1$  is hydrogen or hydroxyl, and  $\text{R}^1$  and  $\text{R}^2$  are unsubstituted  $\text{C}_4$ - $\text{C}_6$  alkoxy, phenoxy or phenyl- $\text{C}_1$ - $\text{C}_4$  alkoxy;

which process comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or alkylation of the phosphonate with the appropriate alkyl halide; or hydrolysis of the diethers or diamines.

2. The process of claim 1 for producing a compound of claim 1 which has the general structural formula II




FORMULA II

wherein

B, R<sup>1</sup> and R<sup>2</sup> are as described in claim 1, provided that when Q is NCHN(R<sup>5</sup>)<sub>2</sub>, then R<sup>5</sup> is not CH<sub>3</sub>;  
X represents hydrogen, CH<sub>2</sub>OR<sup>6</sup> (R,S; or RS stereochemistry), or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> and R<sup>2</sup>, may additionally be independently chosen from OH; and  
R<sup>6</sup> is a hydrolyzable ester group.

3. A process for producing a compound having the general structural formula III




FORMULA III

wherein

B, and R<sup>1</sup> are as previously described in claim 1;  
X represents hydrogen, CH<sub>2</sub>OR<sup>6</sup> (R,S; or R S stereochemistry) or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> may additionally be OH; and R<sup>6</sup> is a hydrolyzable ester group;  
R<sup>7</sup> represents OH, NH<sub>2</sub>, NHR<sup>5</sup>, or NR<sup>5</sup><sub>2</sub>; and  
R<sup>5</sup> is as described in claim 1;  
pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;  
provided that those compounds of the above formula III are excluded wherein  
B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil,  
X is alk<sub>2</sub>-Q<sup>1</sup> wherein is alk<sub>2</sub> selected from a chemical bond or C<sub>1</sub>-C<sub>4</sub> alkylene, Q<sup>1</sup> is hydrogen or hydroxyl, and  
R<sup>1</sup> and R<sup>7</sup> are OH, unsubstituted C<sub>4</sub>-C<sub>6</sub> alkoxy, phenoxy or phenyl-C<sub>1</sub>-C<sub>4</sub> alkoxy;  
which process comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or hydrolysis of the diethers or diamines.

4. A process for producing a compound of claim 1 having the general structural formula IV






FORMULA IV

wherein

$R^8$  and  $R^9$  are identical or different and independently of one another are each  $NR^{12}$ , or oxygen;  
 $R^{10}$  and  $R^{11}$  are identical or different and independently of one another are each hydrogen, or  $R^5$ ;  
 $R^{12}$  represents hydrogen or a lower alkyl;  
 $m$  and  $n$  are identical or different and independently of one another are each 0 or 1;  
 $B$  and  $R^5$  are as described in claim 1; and  
 $X$  is as described in claim 2;

which process comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol.

5. A process for producing a compound having the general structural formula V.



\* stereochemistry is R, S, or RS

FORMULA V

wherein

$R^{13}$  represents  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$ , or OH, provided that  $R^{13}$  is not OH when  $B$  is A or C; and  
 $B$ ,  $R^4$ , and  $R^5$  are as described in claim 1; pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;  
 which process comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or alkylation of the phosphonate with the appropriate alkyl halide.

6. The process of claim 1 for producing a compound of claim 1, wherein  $R^4$  is  $C_4$ - $C_{20}$  alkyl, aryl or aryl-alkyl which may be unsubstituted or substituted by substituents independently selected from hydroxy and halogen.
7. The process of claim 1 for producing a compound of claim 1 wherein  $R^1$  and  $R^2$  are  $OR^4$ , and  $R^4$  represents a physiologically hydrolyzable ester group selected from  $CH_2OC(O)R^5$ ; and  $CH(R^5)OC(O)R^5$  (R,S, or R,S stereochemis-

try); or aryl wherein aryl is substituted by substituents independently selected from hydroxy and halogen.

8. The process of claim 1 for producing a compound of claim 1 which is

5       PMEA, di-(propionyloxymethyl ester);  
       PMEA, di-(isobutyryloxymethyl ester);  
       PMEA, bis-diethylamide;  
       PMEA, di-(butylacetate ester);  
       PMEA, di-(ethylacetate ester);  
 10       PMEA, di-(benzyloxymethyl ester);  
       PMEA, dibutylamide;  
       PMEA, di-(2-methylpropyl ester); or  
       PMEA, di-(3-methylbutyl ester).

15    9. The process of claim 1 for producing a compound of claim 1 which is

      PMEA, di-(pivaloyloxymethyl ester).

10. The process of claim 3 for producing a compound of claim 3 which is

20       PMEA, mono-(pivaloyloxymethyl ester);  
       PMEA, mono-(octyl ester);  
       PMEA, mono-(hydroxy-2,2-dimethylpropyl ester);  
       PMEA, mono-(2,2,2-trichloroethyl ester); or  
 25       PMEA, mono-(2,2-difluoro-3-hydroxypropyl ester).

11. The process of claim 4 for producing a compound of claim 4 which is

      PMEA, cyclic propanyl diester;  
 30       PMEA, cyclic (2,2-dimethyl)propanyl diester;  
       PMEA, cyclic (2,2-dimethyl)propanyl diamide;  
       PMEA, N,N'-dimethyl-cyclic propanyl diamide.

12. The process of claim 1 or claim 3 for producing a compound which is

35       PMEA, di-(isopropyl ester);  
       PMEHx, mono-(isopropyl ester);  
       6-chloro-9-(2-phosphonylmethoxy)ethylpurine,  
       di-(isopropyl ester);  
 40       9-(2-phosphonylmethoxy)ethylpurine, di-(isopropyl ester);  
       PMEA, mono-(isopropyl ester);  
       PMEA, (mono-isopropyl, mono-pivaloyloxymethyl) ester; or  
       PMEA, (mono-isopropyl, mono-phenyl) ester;  
       pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

45    13. The process of claim 1 or claim 3 for producing a compound which is


      PMEA, di-(phenyl ester);  
       PMEA, di-(p-nitrobenzyl ester);  
 50       PMEA, di-(2,2,2-trichloroethyl ester);  
       PMEA, di-(p-trifluoromethylbenzyl ester);  
       PMEA, mono-(choline ester);  
       PMEA, (mono-N,N-diethylacetamide, mono-pivaloyloxymethyl) ester;  
       PMEA, mono-(3-hydroxypropanyl ester);  
 55       PMEA, mono-(3-amino-2,2-dimethylpropyl amide);  
       PMEA, mono-(N,N-diethylacetamide ester);  
       PMEA, mono-(acetic acid ester);  
       PMEA, mono-(N,N-diisopropylacetamide ester);

PMEA, mono-(p-nitrobenzyl ester); or  
 PMEAs, mono-(p-trifluoromethylbenzyl ester);  
 PMEAs, mono-(phenyl ester);  
 pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

14. The use of a compound as defined in anyone of claims 1 to 13 for preparing a pharmaceutical composition for the treatment of viral infection in a mammal.
15. The use of a compound as defined in anyone of claims 1 to 13 for preparing a pharmaceutical composition for inhibiting growth of a tumor in a mammal.
16. A process for preparing a pharmaceutical composition which comprises mixing an amount of at least one compound as defined in anyone of claims 1 to 13 with a pharmaceutically acceptable substantially nontoxic carrier or excipient.

#### Claims for the following Contracting State : GR


1. A compound having the structural formula I



FORMULA I

wherein

B represents adenine (A), cytosine (C), guanine (G), thymine (T), uracil (U), 2,6-diamino purine (DAP), hypoxanthine (Hx),



wherein

Q is independently chosen from H, Cl,  $\text{NHR}^5$ ,  $\text{HR}^5$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH or  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  and  $\text{R}^2$  are identical or different and independently of one another are each  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  or  $\text{N(R}^5)_2$ ;  $\text{R}^1$  and  $\text{R}^2$  optionally being linked with each other to form a cyclic group, or  $\text{R}^1$  or  $\text{R}^2$  optionally being linked to  $\text{R}^3$  to form a cyclic group;

$\text{R}^3$  represents  $\text{C}_1$ - $\text{C}_{20}$  alkylene which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen; or  $\text{R}^3$  is  $\text{CH(CH}_2\text{OR}^5)_2$ , whereby  $\text{R}^1$  and  $\text{R}^2$  each independently may additionally represent OH, and  $\text{R}^5$  is a hydrolyzable ester group;


$\text{R}^4$  represents a physiologically hydrolyzable ester group selected from  $\text{CH}_2\text{C(O)NR}^5$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5\text{)OC(O)R}^5$  (R, S; or RS stereochemistry),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$ , or  $\text{CH}_2\text{OR}^5$ ; or  $\text{R}^4$  represents  $\text{C}_4$ - $\text{C}_{20}$  alkyl, aryl-alkyl or aryl which may be unsubstituted or substituted by substituents independently selected from the group consisting of hydroxy and halogen provided that  $\text{R}^1$  and  $\text{R}^2$  are not simultaneously alkoxy;

$\text{R}^5$  represents  $\text{C}_1$ - $\text{C}_{20}$  alkyl, aryl or aryl-alkyl which may be substituted or unsubstituted by substituents inde-

pendently selected from the group consisting of hydroxy and halogen; pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;


provided that those compounds of the above formula I are excluded wherein

B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil, R<sup>3</sup> is



wherein alk<sub>1</sub> is bonded to B, alk<sub>1</sub>, alk<sub>2</sub> and alk<sub>3</sub> are independently selected from a chemical bond or C<sub>1</sub>-C<sub>4</sub> alkylene, R<sup>a</sup> is hydrogen or C<sub>1</sub>-C<sub>4</sub> alkyl and Q<sup>1</sup> is hydrogen or hydroxyl, and R<sup>1</sup> and R<sup>2</sup> are unsubstituted C<sub>4</sub>-C<sub>6</sub> alkoxy, phenoxy or phenyl-C<sub>1</sub>-C<sub>4</sub> alkoxy.

2. The compound of claim 1 which has the general structural formula II



FORMULA II


wherein

B, R<sup>1</sup> and R<sup>2</sup> are as described in claim 1, provided that when Q is NCHN(R<sup>5</sup>)<sub>2</sub>, then R<sup>5</sup> is not CH<sub>3</sub>;

X represents hydrogen, CH<sub>2</sub>OR<sup>6</sup> (R;S; or RS stereochemistry), or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> and R<sup>2</sup>, may additionally be independently chosen from OH; and

R<sup>6</sup> is a hydrolyzable ester group.

3. A compound having the general structural formula III



FORMULA III

wherein

B, and R<sup>1</sup> are as previously described in claim 1;

X represents hydrogen, CH<sub>2</sub>OR<sup>6</sup> (R;S; or R S stereochemistry) or substituted or unsubstituted lower alkyl, in particular methyl or hydroxymethyl; when X is CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> may additionally be OH; and R<sup>6</sup> is a hydrolyzable ester group;

R<sup>7</sup> represents OH, NH<sub>2</sub>, NHR<sup>5</sup>, or NR<sup>5</sup><sub>2</sub>; and

R<sup>5</sup> is as described in claim 1;


pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof;  
provided that those compounds of the above formula III are excluded wherein

B is adenine, xanthine, hypoxanthine, guanine, 2-aminopurine, 2,6-diaminopurine, cytosine, thymine, or uracil,

X is alk<sub>2</sub>-Q<sup>1</sup> wherein alk<sub>2</sub> is selected from a chemical bond or C<sub>1</sub>-C<sub>4</sub> alkylene, Q<sup>1</sup> is hydrogen or hydroxyl, and

R<sup>1</sup> and R<sup>7</sup> are OH, unsubstituted C<sub>4</sub>-C<sub>6</sub> alkoxy, phenoxy or phenyl-C<sub>1</sub>-C<sub>4</sub> alkoxy.

4. The compound of claim 1 which has the general structural formula IV



FORMULA IV

wherein

R<sup>8</sup> and R<sup>9</sup> are identical or different and independently of one another are each NR<sup>12</sup>, or oxygen;

R<sup>10</sup> and R<sup>11</sup> are identical or different and independently of one another are each hydrogen, or R<sup>5</sup>;


R<sup>12</sup> represents hydrogen or a lower alkyl;

m and n are identical or different and independently of one another are each 0 or 1;

B and R<sup>5</sup> are as described in claim 1; and

X is as described in claim 2.

5. The compound which has the general structural formula V



stereochemistry is R; S; or RS

FORMULA V

wherein

R<sup>13</sup> represents OR<sup>4</sup>, NHR<sup>5</sup>, NR<sup>5</sup>, or OH, provided that R<sup>13</sup> is not OH when B is A or C; and

B, R<sup>4</sup>, and R<sup>5</sup> are as described in claim 1;

pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

6. The compound of claim 1, wherein R<sup>4</sup> is C<sub>4</sub>-C<sub>20</sub> alkyl, aryl or aryl-alkyl which may be unsubstituted or substituted by substituents independently selected from hydroxy and halogen.
7. The compound of claim 1 wherein R<sup>1</sup> and R<sup>2</sup> are OR<sup>4</sup>, and R<sup>4</sup> represents a physiologically hydrolyzable ester group selected from CH<sub>2</sub>OC(O)R<sup>5</sup>; and CH(R<sup>5</sup>)OC(O)R<sup>5</sup> (R;S; or R S stereochemistry); or aryl wherein aryl is substituted by substituents independently selected from hydroxy and halogen.
8. The compound of claim 1 which is

PMEA, di-(propionyloxymethyl ester);  
 PMEA, di-(isobutyryloxymethyl ester);  
 PMEA, bis-diethylamide;  
 PMEA, di-(butylacetate ester);  
 5 PMEA, di-(ethylacetate ester);  
 PMEA, di-(benzoyloxymethyl ester);  
 PMEA, dibutylamide;  
 PMEA, di-(2-methylpropyl ester); or  
 PMEA, di-(3-methylbutyl ester).

10

9. The compound of claim 1 which is

PMEA, di-(pivaloyloxymethyl ester).

15

10. The compound of claim 3 which is

PMEA, mono-(pivaloyloxymethyl ester);  
 PMEA, mono-(octyl ester);  
 PMEA, mono-(hydroxy-2,2-dimethylpropyl ester);  
 20 PMEA, mono-(2,2,2-trichloroethyl ester); or  
 PMEA, mono-(2,2-difluoro-3-hydroxypropyl ester).

11. The compound of claim 4 which is

25

PMEA, cyclic propanyl diester;  
 PMEA, cyclic (2,2-dimethyl)propanyl diester;  
 PMEA, cyclic (2,2-dimethyl)propanyl diamide;  
 PMEA, N,N'-dimethyl-cyclic propanyl diamide.

30

12. The compound which is

PMEA, di-(isopropyl ester);  
 PMEHx, mono-(isopropyl ester);  
 6-chloro-9-(2-phosphonylmethoxy)ethylpurine,  
 35 di-(isopropyl ester);  
 9-(2-phosphonylmethoxy)ethylpurine, di-(isopropyl ester);  
 PMEA, mono-(isopropyl ester);  
 PMEA, (mono-isopropyl, mono-pivaloyloxymethyl) ester; or  
 PMEA, (mono-isopropyl, mono-phenyl) ester;  
 40 pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

13. The compound which is


PMEA, di-(phenyl ester);  
 45 PMEA, di-(p-nitrobenzyl ester);  
 PMEA, di-(2,2,2-trichloroethyl ester);  
 PMEA, di-(p-trifluoromethylbenzyl ester);  
 PMEA, mono-(choline ester);  
 PMEA, (mono-N,N-diethylacetamide, mono-pivaloyloxymethyl) ester;  
 50 PMEA, mono-(3-hydroxypropanyl ester);  
 PMEA, mono-(3-amino-2,2-dimethylpropyl amide);  
 PMEA, mono-(N,N-diethylacetamide ester);  
 PMEA, mono-(acetic acid ester);  
 PMEA, mono-(N,N-diisopropylacetamide ester);  
 55 PMEA, mono-(p-nitrobenzyl ester); or  
 PMEA, mono-(p-trifluoromethylbenzyl ester);  
 PMEA, mono-(phenyl ester);  
 pharmaceutically acceptable acid addition salts, metal salts, and solvates thereof.

14. A process for producing the compound of claim 1, 2, 12, or 13 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or alkylation of the phosphonate with the appropriate alkyl halide; or hydrolysis of the diethers or diamines.
15. A process for producing the compound of claim 3, 12, or 13 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or hydrolysis of the diethers or diamines.
16. A process for producing the compound of claim 4 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol.
17. A process for producing the compound of claim 5 which comprises reacting the phosphonate with an activating agent, then reacting with the appropriate amine or alcohol; or alkylation of the phosphonate with the appropriate alkyl halide.
18. The use of a compound as defined in anyone of claims 1 to 13 for preparing a pharmaceutical composition for the treatment of viral infection in a mammal.
19. The use of a compound as defined in anyone of claims 1 to 13 for preparing a pharmaceutical composition for inhibiting growth of a tumor in a mammal.
20. A process for preparing a pharmaceutical composition which comprises mixing an amount of at least one compound as defined in anyone of claims 1 to 13 with a pharmaceutically acceptable substantially nontoxic carrier or excipient.

#### Patentansprüche


Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE

#### 1. Verbindung mit der Strukturformel I




worin

B für Adenin (A), Cytosin (C), Guanin (G), Thymin (T), Uracil (U), 2,6-Diaminopurin (DAP), Hypoxanthin (Hx) steht,



oder



worin

Q unabhängig ausgewählt ist aus H, Cl, NHR<sup>5</sup>, NR<sup>5</sup><sub>2</sub>, NHC(O)R<sup>5</sup>, N(C(O)R<sup>5</sup>)<sub>2</sub>, OH oder NCHN(R<sup>5</sup>)<sub>2</sub>;

R<sup>1</sup> und R<sup>2</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für OR<sup>4</sup>, NH<sub>2</sub>, NHR<sup>5</sup> oder N(R<sup>5</sup>)<sub>2</sub> stehen; R<sup>1</sup> und R<sup>2</sup> gegebenenfalls miteinander unter Bildung einer cyclischen Gruppe verbunden sind oder R<sup>1</sup> oder R<sup>2</sup> gegebenenfalls mit R<sup>3</sup> unter Bildung einer cyclischen Gruppe verbunden sind;

R<sup>3</sup> für C<sub>1</sub>-C<sub>20</sub>-Alkylen steht, das unsubstituiert oder mit unabhängig aus Hydroxy und Halogen ausgewählten

Substituenten substituiert ist; oder  $R^3$  für  $\text{CH}(\text{CH}_2\text{OR}^6)\text{CH}_2$  steht, wobei  $R^1$  und  $R^2$  jeweils unabhängig zusätzlich für OH stehen können und  $R^6$  eine hydrolysierbare Estergruppe ist;


$R^4$  für eine physiologisch hydrolysierbare Estergruppe steht, die ausgewählt ist unter  $\text{CH}_2\text{C}(\text{O})\text{NR}^5_2$ ,  $\text{CH}_2\text{C}(\text{O})\text{OR}^5$ ,  $\text{CH}_2\text{OC}(\text{O})\text{R}^5$ ,  $\text{CH}(\text{R}^5)\text{OC}(\text{O})\text{R}^5$  ( $R$ : S; oder RS Stereochemie),  $\text{CH}_2\text{C}(\text{R}^5)_2\text{CH}_2\text{OH}$  oder  $\text{CH}_2\text{OR}^5$ ; oder  $R^4$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $\text{C}_4$ - $\text{C}_{20}$ -Alkyl, Aryl-Alkyl oder Aryl steht, mit der Maßgabe, daß  $R^1$  und  $R^2$  nicht gleichzeitig Alkoxy sind;

$R^5$  für unsubstituierte oder durch unabhängig aus Hydroxy und Halogen ausgewählten Substituenten substituiertes  $\text{C}_1$ - $\text{C}_{20}$ -Alkyl, Aryl oder Aryl-Alkyl steht;

pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon;


mit der Maßgabe, daß diejenigen Verbindungen obiger Formel I ausgeschlossen sind, worin

B für Adenin, Xanthin, Hypoxanthin, Guanin, 2-Aminopurin, 2,6-Diaminopurin, Cytosin, Thymin oder Uracil steht,  $R^3$  für



steht worin  $\text{alk}_1$  an B gebunden ist,  $\text{alk}_1$ ,  $\text{alk}_2$  und  $\text{alk}_3$  unabhängig voneinander ausgewählt sind unter einer chemischen Bindung oder  $\text{C}_1$ - $\text{C}_4$ -Alkylen,  $R^a$  für Wasserstoff oder  $\text{C}_1$ - $\text{C}_4$ -Alkyl und  $Q^1$  für Wasserstoff oder Hydroxyl steht und  $R^1$  und  $R^2$  für unsubstituiertes  $\text{C}_4$ - $\text{C}_6$ -Alkoxy, Phenoxy oder Phenyl- $\text{C}_1$ - $\text{C}_4$ -Alkoxy stehen.

## 2. Verbindung nach Anspruch 1 mit der allgemeinen Strukturformel II



Formel II

worin


B,  $R^1$  und  $R^2$  wie in Anspruch 1 definiert sind, mit der Maßgabe, daß  $R^5$  nicht  $\text{CH}_3$  ist, falls Q für  $\text{NCHN}(\text{R}^5)_2$  steht;

X für Wasserstoff,  $\text{CH}_2\text{OR}^6$  ( $R$ : S; oder RS Stereochemie), oder substituiertes oder unsubstituiertes Niedrigalkyl, insbesondere Methyl oder Hydroxymethyl steht; falls X für  $\text{CH}_2\text{OR}^6$  steht,  $R^1$  und  $R^2$  unabhängig zusätzlich OH sein können; und

$R^6$  für eine hydrolysierbare Estergruppe steht.

## 3. Verbindung mit der allgemeinen Strukturformel III





Formel III

10 worin

B und R<sup>1</sup> wie zuvor in Anspruch 1 definiert sind;

X für Wasserstoff, CH<sub>2</sub>OR<sup>6</sup> (R; S; oder RS Stereochemie), oder substituiertes oder unsubstituiertes Niedrigalkyl, insbesondere Methyl oder Hydroxymethyl steht; falls X für CH<sub>2</sub>OR<sup>6</sup> steht, R<sup>1</sup> zusätzlich OH sein kann; und R<sup>6</sup> eine hydrolysierbare Estergruppe ist;

R<sup>7</sup> für OH, NH<sub>2</sub>, NHR<sup>5</sup> oder NR<sup>5</sup><sub>2</sub> steht; und


R<sup>5</sup> wie in Anspruch 1 definiert ist; pharmazeutische akzeptable Säureadditionssalze, Metallsalze und Solvate davon;

mit der Maßgabe, daß diejenigen Verbindungen obiger Formel III ausgeschlossen sind, worin

B für Adenin, Xanthin, Hypoxanthin, Guanin, 2-Aminopurin, 2,6-Diaminopurin, Cytosin, Thymin oder Uracil steht,

X für alk<sub>2</sub>-Q<sup>1</sup> steht, worin alk<sub>2</sub> ausgewählt ist unter einer chemischen Bindung oder C<sub>1</sub>-C<sub>4</sub>-Alkylen, Q<sup>1</sup> für Wasserstoff oder Hydroxyl steht und R<sup>1</sup> und R<sup>7</sup> für OH, unsubstituiertes C<sub>4</sub>-C<sub>6</sub>-Alkoxy, Phenoxy oder Phenyl-C<sub>1</sub>-C<sub>4</sub>-Alkoxy stehen.

4. Verbindung nach Anspruch 1 mit der allgemeinen Strukturformel IV



Formel IV

worin

R<sup>8</sup> und R<sup>9</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für NR<sup>12</sup> oder Sauerstoff stehen; R<sup>10</sup> und R<sup>11</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für Wasserstoff oder R<sup>5</sup> stehen;


R<sup>12</sup> für Wasserstoff oder Niedrigalkyl steht;

m und n gleich oder verschieden sind und unabhängig voneinander jeweils für 0 oder 1 stehen;

B und R<sup>5</sup> wie in Anspruch 1 definiert sind; und

X wie in Anspruch 2 definiert ist.

5. Verbindung mit der allgemeinen Strukturformel V



Formel V

Stereochemie ist R; S; oder RS

worin

$R^{13}$  für  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$  oder OH steht, mit der Maßgabe, daß  $R^{13}$  nicht OH ist, falls B für A oder C steht; und B,  $R^4$ , und  $R^5$  wie in Anspruch 1 definiert sind;

pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon.

6. Verbindung nach Anspruch 1, worin  $R^4$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $C_4$ - $C_{20}$ -Alkyl, Aryl oder Aryl-Alkyl steht.

7. Verbindung nach Anspruch 1, worin  $R^1$  und  $R^2$  für  $OR^4$  und  $R^4$  für eine aus  $CH_2OC(O)R^5$  und  $CH(R^5)OC(O)R^5$  ( $R$ ; S; oder RS Stereochemie) ausgewählte, physiologisch hydrolysierbare Estergruppe oder Aryl steht, worin Aryl mit unabhängig aus Hydroxy und Halogen ausgewählten Substituenten substituiert ist.

8. Verbindung nach Anspruch 1, nämlich

PMEA, Di(propionyloxymethylester);

PMEA, Di(isobutyryloxymethylester);

PMEA, Bis(diethylamid);

PMEA, Di(butylacetatester);

PMEA, Di(ethylacetatester);

PMEA, Di(benzoyloxymethylester);

PMEA, Dibutylamid;

PMEA, Di(2-methylpropylester); oder

PMEA, Di(3-methylbutylester).

9. Verbindung nach Anspruch 1, nämlich

PMEA, Di(pivaloyloxymethylester).

10. Die Verbindung nach Anspruch 3, nämlich

PMEA, Mono(pivaloyloxymethylester);

PMEA, Mono(octylester);

PMEA, Mono(hydroxy-2,2-dimethylpropylester);

PMEA, Mono-(2,2,2-trichlorethylester); oder

PMEA, Mono-(2,2-difluor-3-hydroxypropylester).

11. Verbindung nach Anspruch 4, nämlich

PMEA, cyclischer Propanyldiester;

PMEA, cyclischer (2,2-Dimethyl)propanyldiester;

PMEA, cyclisches (2,2-Dimethyl)propanyldiamid;

PMEA, N,N'-Dimethyl-cyclisches Propanyldiamid.

12. Verbindung, nämlich

PMEA, Di(isopropylester);

PMEHx, Mono(isopropylester);

6-Chlor-9-(2-phosphonylmethoxy)ethylpurin,

Di(isopropylester);

9-(2-Phosphonylmethoxy)ethylpurin, Di(isopropylester);

PMEA, Mono(isopropylester);

PMEA, (Monoisopropyl, monopivaloyloxymethyl)ester; oder

PMEA, (Monoisopropyl, monophenyl)ester;

pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon.


13. Verbindung, nämlich

PMEA, Di(phenylester);  
 PMEA, Di-(p-nitrobenzylester);  
 PMEA, Di-(2,2,2-trichlorethylester);  
 PMEA, Di-(p-trifluormethylbenzylester);  
 5 PMEA, Mono(cholinester);  
 PMEA, (Mono-N,N-diethylacetamid, monopivaloyloxymethyl)  
 ester;  
 PMEA, Mono-(3-hydroxypropanylester);  
 PMEA, Mono-(3-amino-2,2-dimethylpropylamid);  
 10 PMEA, Mono-(N,N-diethylacetamidester);  
 PMEA, Mono(essigsäureester);  
 PMEA, Mono-(N,N-diisopropylacetamidester);  
 PMEA, Mono-(p-nitrobenzylester); oder  
 PMEA, Mono-(p-trifluormethylbenzylester);  
 15 PMEA, Mono(phenylester);  
 pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon.

14. Verfahren zur Herstellung einer Verbindung nach Anspruch 1, 2, 12 oder 13, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder das Phosphonat mit dem geeigneten Alkylhalogenid alkylert; oder die Diether oder Diamine hydrolysiert.
15. Verfahren zur Herstellung einer Verbindung nach Anspruch 3, 12 oder 13, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder die Diether oder Diamine hydrolysiert.
16. Verfahren zur Herstellung einer Verbindung nach Anspruch 4, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt.
17. Verfahren zur Herstellung einer Verbindung nach Anspruch 5, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder das Phosphonat mit dem geeigneten Alkylhalogenid alkylert.
18. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 13 zur Herstellung einer pharmazeutischen Zusammensetzung zur Behandlung viraler Infektionen eines Säugers.
19. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 13 zur Herstellung einer pharmazeutischen Zusammensetzung zur Wachstums hemmung eines Säugertumors.
20. Pharmazeutische Zusammensetzung mit wenigstens einer Verbindung der Ansprüche 1 bis 13 zusammen mit einem pharmazeutisch akzeptablen, im wesentlichen atoxischen Träger oder Excipienten.


#### Patentansprüche für folgenden Vertragsstaat : ES

1. Verfahren zur Herstellung einer Verbindung mit der Strukturformel I




worin

B für Adenin (A), Cytosin (C), Guanin (G), Thymin (T), Uracil (U), 2,6-Diaminopurin (DAP), Hypoxanthin (Hx) steht,



oder



worin

Q unabhängig ausgewählt ist aus H, Cl,  $\text{NHR}^5$ ,  $\text{NR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH oder  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  und  $\text{R}^2$  gleich oder verschieden sind und unabhängig voneinander jeweils für  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  oder  $\text{N(R}^5)_2$  stehen;  $\text{R}^1$  und  $\text{R}^2$  gegebenenfalls miteinander unter Bildung einer cyclischen Gruppe verbunden sind oder  $\text{R}^1$  oder  $\text{R}^2$  gegebenenfalls mit  $\text{R}^3$  unter Bildung einer cyclischen Gruppe verbunden sind;

$\text{R}^3$  für  $\text{C}_1\text{-C}_{20}$ -Alkylen steht, das unsubstituiert oder mit unabhängig aus Hydroxy und Halogen ausgewählten Substituenten substituiert ist; oder  $\text{R}^3$  für  $\text{CH(CH}_2\text{OR}^6)_2$  steht, wobei  $\text{R}^1$  und  $\text{R}^2$  jeweils unabhängig zusätzlich für OH stehen können und  $\text{R}^6$  eine hydrolysierbare Estergruppe ist;


$\text{R}^4$  für eine physiologisch hydrolysierbare Estergruppe steht, die ausgewählt ist unter  $\text{CH}_2\text{C(O)NR}^5_2$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5)_2\text{OC(O)R}^5$  ( $\text{R}^5$ : S; oder RS Stereochemie),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$  oder  $\text{CH}_2\text{OR}^5$ ; oder  $\text{R}^4$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $\text{C}_4\text{-C}_{20}$ -Alkyl, Aryl-Alkyl oder Aryl steht, mit der Maßgabe, daß  $\text{R}^1$  und  $\text{R}^2$  nicht gleichzeitig Alkoxy sind;

$\text{R}^5$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $\text{C}_1\text{-C}_{20}$ -Alkyl, Aryl oder Aryl-Alkyl steht;

pharmazeutisch akzeptabler Säureadditionssalze, Metallsalze und Solvate davon;


mit der Maßgabe, daß diejenigen Verbindungen obiger Formel I ausgeschlossen sind, worin

B für Adenin, Xanthin, Hypoxanthin, Guanin, 2-Aminopurin, 2,6-Diaminopurin, Cytosin, Thymin oder Uracil steht,  $\text{R}^3$  für



steht worin  $\text{alk}_1$  an B gebunden ist,  $\text{alk}_1$ ,  $\text{alk}_2$  und  $\text{alk}_3$  unabhängig voneinander ausgewählt sind unter einer chemischen Bindung oder  $\text{C}_1\text{-C}_4$ -Alkylen,  $\text{R}^a$  für Wasserstoff oder  $\text{C}_1\text{-C}_4$ -Alkyl und  $\text{Q}^1$  für Wasserstoff oder Hydroxyl steht und  $\text{R}^1$  und  $\text{R}^2$  für unsubstituiertes  $\text{C}_4\text{-C}_6$ -Alkoxy, Phenoxy oder Phenyl- $\text{C}_1\text{-C}_4$ -Alkoxy stehen; wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder das Phosphonat mit dem geeigneten Alkylhalogenid alkyliert; oder die Diether oder Diamine hydrolysiert.

## 2. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung aus Anspruch 1 mit der allgemeinen Strukturformel II



Formel II


worin

B, R<sup>1</sup> und R<sup>2</sup> wie in Anspruch 1 definiert sind, mit der Maßgabe, daß R<sup>5</sup> nicht CH<sub>3</sub> ist, falls Q für NCHN(R<sup>5</sup>)<sub>2</sub> steht;

X für Wasserstoff, CH<sub>2</sub>OR<sup>6</sup> (R; S; oder RS Stereochemie), oder substituiertes oder unsubstituiertes Niedrigalkyl, insbesondere Methyl oder Hydroxymethyl steht; falls X für CH<sub>2</sub>OR<sup>6</sup> steht, R<sup>1</sup> und R<sup>2</sup> unabhängig zusätzlich OH sein können; und

R<sup>6</sup> für eine hydrolysierbare Estergruppe steht.

3. Verfahren zur Herstellung einer Verbindung mit der allgemeinen Strukturformel III



Formel III

worin

B und R<sup>1</sup> wie zuvor in Anspruch 1 definiert sind;

X für Wasserstoff, CH<sub>2</sub>OR<sup>6</sup> (R; S; oder RS Stereochemie), oder substituiertes oder unsubstituiertes Niedrigalkyl, insbesondere Methyl oder Hydroxymethyl steht; falls X für CH<sub>2</sub>OR<sup>6</sup> steht, R<sup>1</sup> zusätzlich OH sein kann; und R<sup>6</sup> eine hydrolysierbare Estergruppe ist;

R<sup>7</sup> für OH, NH<sub>2</sub>, NHR<sup>5</sup> oder NR<sup>5</sup><sub>2</sub> steht; und

R<sup>5</sup> wie in Anspruch 1 definiert ist; pharmazeutische akzeptabler Säureadditionssalze, Metallsalze und Solvate davon;


mit der Maßgabe, daß diejenigen Verbindungen obiger Formel III ausgeschlossen sind, worin

B für Adenin, Xanthin, Hypoxanthin, Guanin, 2-Aminopurin, 2,6-Diaminopurin, Cytosin, Thymin oder Uracil steht,

X für alk<sub>2</sub>-Q<sup>1</sup> steht, worin alk<sub>2</sub> ausgewählt ist unter einer chemischen Bindung oder C<sub>1</sub>-C<sub>4</sub>-Alkylen, Q<sup>1</sup> für Wasserstoff oder Hydroxyl steht und R<sup>1</sup> und R<sup>7</sup> für OH, unsubstituiertes C<sub>4</sub>-C<sub>6</sub>-Alkoxy, Phenoxy oder Phenyl-C<sub>1</sub>-C<sub>4</sub>-Alkoxy stehen;

wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder die Diether oder Diamine hydrolysiert.

4. Verfahren zur Herstellung einer Verbindung aus Anspruch 1 mit der allgemeinen Strukturformel IV



Formel IV

worin

R<sup>8</sup> und R<sup>9</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für NR<sup>12</sup> oder Sauerstoff stehen; R<sup>10</sup> und R<sup>11</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für Wasserstoff oder R<sup>5</sup> stehen;


R<sup>12</sup> für Wasserstoff oder Niedrigalkyl steht;

m und n gleich oder verschieden sind und unabhängig voneinander jeweils für 0 oder 1 stehen;

B und R<sup>5</sup> wie in Anspruch 1 definiert sind; und

X wie in Anspruch 2 definiert ist;  
wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt.

- 5 5. Verfahren zur Herstellung einer Verbindung mit der allgemeinen Strukturformel V



Formel V

Stereochemie ist R; S; oder RS

15 worin

20  $R^{13}$  für  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$  oder OH steht, mit der Maßgabe, daß  $R^{13}$  nicht OH ist, falls B für A oder C steht; und B,  $R^4$ , und  $R^5$  wie in Anspruch 1 definiert sind; pharmazeutisch akzeptabler Säureadditionssalze, Metallsalze und Solvate davon;  
wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder das Phosphonat mit dem geeigneten Alkylhalogenid alkyliert.

- 25 6. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung aus Anspruch 1, worin  $R^4$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $C_4$ - $C_{20}$ -Alkyl, Aryl oder Aryl-Alkyl steht.

- 30 7. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung aus Anspruch 1, worin  $R^1$  und  $R^2$  für  $OR^4$  und  $R^4$  für eine aus  $CH_2OC(O)R^5$  und  $CH(R^5)OC(O)R^5$  ( $R$ ; S; oder RS Stereochemie) ausgewählte, physiologisch hydrolysierbare Estergruppe oder Aryl steht, worin Aryl mit unabhängig aus Hydroxy und Halogen ausgewählten Substituenten substituiert ist.

8. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung aus Anspruch 1, nämlich

35 PMEA, Di(propionyloxymethylester);  
PMEA, Di(isobutyryloxymethylester);  
PMEA, Bis(diethylamid);  
PMEA, Di(butylacetatester);  
PMEA, Di(ethylacetatester);  
40 PMEA, Di(benzoyloxymethylester);  
PMEA, Dibutylamid;  
PMEA, Di(2-methylpropylester); oder  
PMEA, Di(3-methylbutylester).

- 45 9. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung aus Anspruch 1, nämlich

PMEA, Di(pivaloyloxymethylester).

10. Verfahren nach Anspruch 3 zur Herstellung einer Verbindung aus Anspruch 3, nämlich

50 PMEA, Mono(pivaloyloxymethylester);  
PMEA, Mono(octylester);  
PMEA, Mono(hydroxy-2,2-dimethylpropylester);  
PMEA, Mono-(2,2,2-trichlorethylester); oder  
55 PMEA, Mono-(2,2-difluor-3-hydroxypropylester).

11. Verfahren nach Anspruch 4 zur Herstellung einer Verbindung aus Anspruch 4, nämlich

PMEA, cyclischer Propanyldiester;  
 PMEA, cyclischer (2,2-Dimethyl)propanyldiester;  
 PMEA, cyclisches (2,2-Dimethyl)propanyldiamid;  
 PMEA, N,N'-Dimethyl-cyclisches Propanyldiamid.

12. Verfahren nach Anspruch 1 oder Anspruch 3 zur Herstellung einer Verbindung, nämlich

PMEA, Di(isopropylester);  
 PMEHx, Mono(isopropylester);  
 6-Chlor-9-(2-phosphonylmethoxy)ethylpurin,  
 Di(isopropylester);  
 9-(2-Phosphonylmethoxy)ethylpurin, Di(isopropylester);  
 PMEA, Mono(isopropylester);  
 PMEA, (Monoisopropyl, monopivaloyloxymethyl)ester; oder  
 PMEA, (Monoisopropyl, monophenyl)ester;  
 pharmazeutisch akzeptabler Säureadditionssalze, Metallsalze und Solvate davon.

13. Verfahren nach Anspruch 1 oder Anspruch 3 zur Herstellung einer Verbindung, nämlich

PMEA, Di(phenylester);  
 PMEA, Di-(p-nitrobenzylester);  
 PMEA, Di-(2,2,2-trichlorethylester);  
 PMEA, Di-(p-trifluormethylbenzylester);  
 PMEA, Mono(cholinester);  
 PMEA, (Mono-N,N-diethylacetamid, monopivaloyloxymethyl)ester;  
 PMEA, Mono-(3-hydroxypropylester);  
 PMEA, Mono-(3-amino-2,2-dimethylpropylamid);  
 PMEA, Mono-(N,N-diethylacetamidester);  
 PMEA, Mono(essigsäureester);  
 PMEA, Mono-(N,N-diisopropylacetamidester);  
 PMEA, Mono-(p-nitrobenzylester); oder  
 PMEA, Mono-(p-trifluormethylbenzylester);  
 PMEA, Mono(phenylester);  
 pharmazeutisch akzeptabler Säureadditionssalze, Metallsalze und Solvate davon.


14. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 13 zur Herstellung einer pharmazeutischen Zusammensetzung zur Behandlung viraler Infektionen eines Säugers.

15. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 13 zur Herstellung einer pharmazeutischen Zusammensetzung zur Wachstumshemmung eines Säugertumors.

16. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, wobei man eine Menge wenigstens einer in einem der Ansprüche 1 bis 13 definierten Verbindung mit einem pharmazeutisch akzeptablen, im wesentlichen atoxischen Träger oder Excipienten mischt.


Patentansprüche für folgenden Vertragsstaat : GR

1. Verbindung mit der Strukturformel I



worin

B für Adenin (A), Cytosin (C), Guanin (G), Thymin (T), Uracil (U), 2,6-Diaminopurin (DAP), Hypoxanthin (Hx) steht,



worin

Q unabhängig ausgewählt ist aus H, Cl,  $\text{NHR}^5$ ,  $\text{NR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH oder  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  und  $\text{R}^2$  gleich oder verschieden sind und unabhängig voneinander jeweils für  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  oder  $\text{N(R}^5)_2$  stehen;  $\text{R}^1$  und  $\text{R}^2$  gegebenenfalls miteinander unter Bildung einer cyclischen Gruppe verbunden sind oder  $\text{R}^1$  oder  $\text{R}^2$  gegebenenfalls mit  $\text{R}^3$  unter Bildung einer cyclischen Gruppe verbunden sind;

$\text{R}^3$  für  $\text{C}_1$ - $\text{C}_{20}$ -Alkylen steht, das unsubstituiert oder mit unabhängig aus Hydroxy und Halogen ausgewählten Substituenten substituiert ist; oder  $\text{R}^3$  für  $\text{CH(CH}_2\text{OR}^6)_2$  steht, wobei  $\text{R}^1$  und  $\text{R}^2$  jeweils unabhängig zusätzlich für OH stehen können und  $\text{R}^6$  eine hydrolysierbare Estergruppe ist;


$\text{R}^4$  für eine physiologisch hydrolysierbare Estergruppe steht, die ausgewählt ist unter  $\text{CH}_2\text{C(O)NR}^5_2$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5)_2\text{OC(O)R}^5$  ( $\text{R}$ : S; oder RS Stereochemie),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$  oder  $\text{CH}_2\text{OR}^5$ ; oder  $\text{R}^4$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $\text{C}_4$ - $\text{C}_{20}$ -Alkyl, Aryl-Alkyl oder Aryl steht, mit der Maßgabe, daß  $\text{R}^1$  und  $\text{R}^2$  nicht gleichzeitig Alkoxy sind;

$\text{R}^5$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $\text{C}_1$ - $\text{C}_{20}$ -Alkyl, Aryl oder Aryl-Alkyl steht;

pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon;


mit der Maßgabe, daß diejenigen Verbindungen obiger Formel I ausgeschlossen sind, worin

B für Adenin, Xanthin, Hypoxanthin, Guanin, 2-Aminopurin, 2,6-Diaminopurin, Cytosin, Thymin oder Uracil steht,  $\text{R}^3$  für



steht worin  $\text{alk}_1$  an B gebunden ist,  $\text{alk}_1$ ,  $\text{alk}_2$  und  $\text{alk}_3$  unabhängig voneinander ausgewählt sind unter einer chemischen Bindung oder  $\text{C}_1$ - $\text{C}_4$ -Alkylen,  $\text{R}^a$  für Wasserstoff oder  $\text{C}_1$ - $\text{C}_4$ -Alkyl und  $\text{Q}^1$  für Wasserstoff oder Hydroxyl steht und  $\text{R}^1$  und  $\text{R}^2$  für unsubstituiertes  $\text{C}_4$ - $\text{C}_6$ -Alkoxy, Phenoxy oder Phenyl- $\text{C}_1$ - $\text{C}_4$ -Alkoxy stehen.

## 2. Verbindung nach Anspruch 1 mit der allgemeinen Strukturformel II



worin




B, R<sup>1</sup> und R<sup>2</sup> wie in Anspruch 1 definiert sind, mit der Maßgabe, daß R<sup>5</sup> nicht CH<sub>3</sub> ist, falls Q für NCHN(R<sup>5</sup>)<sub>2</sub> steht;

X für Wasserstoff, CH<sub>2</sub>OR<sup>6</sup> (R; S; oder RS Stereochemie), oder substituiertes oder unsubstituiertes Niedrigalkyl, insbesondere Methyl oder Hydroxymethyl steht; falls X für CH<sub>2</sub>OR<sup>6</sup> steht, R<sup>1</sup> und R<sup>2</sup> unabhängig zusätzlich OH sein können; und

R<sup>6</sup> für eine hydrolysierbare Estergruppe steht.

### 3. Verbindung mit der allgemeinen Strukturformel III



Formel III

worin

B und R<sup>1</sup> wie zuvor in Anspruch 1 definiert sind;

X für Wasserstoff, CH<sub>2</sub>OR<sup>6</sup> (R; S; oder RS Stereochemie), oder substituiertes oder unsubstituiertes Niedrigalkyl, insbesondere Methyl oder Hydroxymethyl steht; falls X für CH<sub>2</sub>OR<sup>6</sup> steht, R<sup>1</sup> zusätzlich OH sein kann; und R<sup>6</sup> eine hydrolysierbare Estergruppe ist;

R<sup>7</sup> für OH, NH<sub>2</sub>, NHR<sup>5</sup> oder NR<sup>5</sup><sub>2</sub> steht; und

R<sup>5</sup> wie in Anspruch 1 definiert ist;


pharmazeutische akzeptable Säureadditionssalze, Metallsalze und Solvate davon;

mit der Maßgabe, daß diejenigen Verbindungen obiger Formel III ausgeschlossen sind, worin

B für Adenin, Xanthin, Hypoxanthin, Guanin, 2-Aminopurin, 2,6-Diaminopurin, Cytosin, Thymin oder Uracil steht,

X für alk<sub>2</sub>-Q<sup>1</sup> steht, worin alk<sub>2</sub> ausgewählt ist unter einer chemischen Bindung oder C<sub>1</sub>-C<sub>4</sub>-Alkylen, Q<sup>1</sup> für Wasserstoff oder Hydroxyl steht und R<sup>1</sup> und R<sup>7</sup> für OH, unsubstituiertes C<sub>4</sub>-C<sub>6</sub>-Alkoxy, Phenoxy oder Phenyl-C<sub>1</sub>-C<sub>4</sub>-Alkoxy stehen.

### 4. Verbindung nach Anspruch 1 mit der allgemeinen Strukturformel IV



Formel IV

worin

R<sup>8</sup> und R<sup>9</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für NR<sup>12</sup> oder Sauerstoff stehen; R<sup>10</sup> und R<sup>11</sup> gleich oder verschieden sind und unabhängig voneinander jeweils für Wasserstoff oder R<sup>5</sup> stehen;


R<sup>12</sup> für Wasserstoff oder Niedrigalkyl steht;

m und n gleich oder verschieden sind und unabhängig voneinander jeweils für 0 oder 1 stehen;

B und R<sup>5</sup> wie in Anspruch 1 definiert sind; und

X wie in Anspruch 2 definiert ist.

### 5. Verbindung mit der allgemeinen Strukturformel V



Formel V

Stereochemie ist R; S; oder RS

worin

$R^{13}$  für  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$  oder OH steht, mit der Maßgabe, daß  $R^{13}$  nicht OH ist, falls B für A oder C steht; und B,  $R^4$ , und  $R^5$  wie in Anspruch 1 definiert sind; pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon.

6. Verbindung nach Anspruch 1, worin  $R^4$  für unsubstituiertes oder durch unabhängig aus Hydroxy und Halogen ausgewählte Substituenten substituiertes  $C_4$ - $C_{20}$ -Alkyl, Aryl oder Aryl-Alkyl steht.

7. Verbindung nach Anspruch 1, worin  $R^1$  und  $R^2$  für  $OR^4$  und  $R^4$  für eine aus  $CH_2OC(O)R^5$  und  $CH(R^5)OC(O)R^5$  ( $R$ ; S; oder RS Stereochemie) ausgewählte, physiologisch hydrolysierbare Estergruppe oder Aryl steht, worin Aryl mit unabhängig aus Hydroxy und Halogen ausgewählten Substituenten substituiert ist.

8. Verbindung nach Anspruch 1, nämlich

PMEA, Di(propionyloxymethylester);  
 PMEa, Di(isobutyryloxymethylester);  
 PMEa, Bis(diethylamid);  
 PMEa, Di(butylacetatester);  
 PMEa, Di(ethylacetatester);  
 PMEa, Di(benzoyloxymethylester);  
 PMEa, Dibutylamid;  
 PMEa, Di(2-methylpropylester); oder  
 PMEa, Di(3-methylbutylester).

9. Verbindung nach Anspruch 1, nämlich

PMEA, Di(pivaloyloxymethylester).

10. Die Verbindung nach Anspruch 3, nämlich

PMEA, Mono(pivaloyloxymethylester);  
 PMEa, Mono(octylester);  
 PMEa, Mono(hydroxy-2,2-dimethylpropylester);  
 PMEa, Mono-(2,2,2-trichlorethylester); oder  
 PMEa, Mono-(2,2-difluor-3-hydroxypropylester).

11. Verbindung nach Anspruch 4, nämlich

PMEA, cyclischer Propanyldiester;  
 PMEa, cyclischer (2,2-Dimethyl)propanyldiester;  
 PMEa, cyclisches (2,2-Dimethyl)propanyldiamid;  
 PMEa, N,N-Dimethyl-cyclisches Propanyldiamid.

12. Verbindung, nämlich

PMEA, Di(isopropylester);

PMEHx, Mono(isopropylester);  
 6-Chlor-9-(2-phosphonylmethoxy)ethylpurin,  
 Di(isopropylester);  
 9-(2-Phosphonylmethoxy)ethylpurin, Di(isopropylester);  
 5   PMEA, Mono(isopropylester);  
 PMEa, (Monoisopropyl, monopivaloyloxymethyl)ester; oder  
 PMEa, (Monoisopropyl, monophenyl)ester;  
 pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon.

10   13. Verbindung, nämlich

PMEA, Di(phenylester);  
 PMEa, Di-(p-nitrobenzylester);  
 PMEa, Di-(2,2,2-trichlorethylester);  
 15   PMEA, Di-(p-trifluormethylbenzylester);  
 PMEa, Mono(cholinester);  
 PMEa, (Mono-N,N-diethylacetamid, monopivaloyloxymethyl)ester;  
 PMEa, Mono-(3-hydroxypropylester);  
 PMEa, Mono-(3-amino-2,2-dimethylpropylamid);  
 20   PMEA, Mono-(N,N-diethylacetamidester);  
 PMEa, Mono(essigsäureester);  
 PMEa, Mono-(N,N-diisopropylacetamidester);  
 PMEa, Mono-(p-nitrobenzylester); oder  
 PMEa, Mono-(p-trifluormethylbenzylester);  
 25   PMEA, Mono(phenylester);  
 pharmazeutisch akzeptable Säureadditionssalze, Metallsalze und Solvate davon.

14. Verfahren zur Herstellung einer Verbindung nach Anspruch 1, 2, 12 oder 13, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder das Phosphonat mit dem geeigneten Alkylhalogenid alkyliert; oder die Diether oder Diamine hydrolysiert.

15. Verfahren zur Herstellung einer Verbindung nach Anspruch 3, 12 oder 13, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder die Diether oder Diamine hydrolysiert.

16. Verfahren zur Herstellung einer Verbindung nach Anspruch 4, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt.

17. Verfahren zur Herstellung einer Verbindung nach Anspruch 5, wobei man das Phosphonat mit einem Aktivierungsmittel und dann mit dem geeigneten Amin oder Alkohol umsetzt; oder das Phosphonat mit dem geeigneten Alkylhalogenid alkyliert.

18. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 13 zur Herstellung einer pharmazeutischen Zusammensetzung zur Behandlung viraler Infektionen eines Säugers.


19. Verwendung einer Verbindung nach einem der Ansprüche 1 bis 13 zur Herstellung einer pharmazeutischen Zusammensetzung zur Wachstumshemmung eines Säugertumors.

20. Verfahren zur Herstellung einer pharmazeutischen Zusammensetzung, wobei man eine Menge wenigstens einer in einem der Ansprüche 1 bis 13 definierten Verbindung mit einem pharmazeutisch akzeptablen, im wesentlichen atoxischen Träger oder Exipienten mischt.

Revendications

55   Revendications pour les Etats contractants suivants : AT, BE, CH, DE, DK, FR, GB, IT, LI, LU, NL, SE


1. Composé ayant la formule développée I :



FORMULE I

dans laquelle

B représente l'adénine (A), la cytosine (C), la guanine (G), la thymine (T), l'uracile (U), la 2,6-diaminopurine (DAP), l'hypoxanthine (Hx);



dans laquelle

Q est indépendamment choisi parmi H, Cl,  $\text{NHR}^5$ ,  $\text{NHR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH ou  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  et  $\text{R}^2$  sont identiques ou différents et représentent indépendamment  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  ou  $\text{N(R}^5)_2$ ;  $\text{R}^1$  et  $\text{R}^2$  étant éventuellement liés l'un à l'autre pour former un groupe cyclique, ou  $\text{R}^1$  ou  $\text{R}^2$  étant éventuellement liés à  $\text{R}^3$  pour former un groupe cyclique;


$\text{R}^3$  représente un groupe alkylène en  $\text{C}_1$  à  $\text{C}_{20}$  qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène; ou  $\text{R}^3$  représente  $\text{CH(CH}_2\text{OR}^6)_2$ , auquel cas  $\text{R}^1$  et  $\text{R}^2$  peuvent en outre représenter indépendamment OH, et  $\text{R}^6$  représente un groupe ester hydrolysable;

$\text{R}^4$  représente un groupe ester physiologiquement hydrolysable choisi parmi  $\text{CH}_2\text{C(O)NR}^5$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5\text{)OC(O)R}^5$  (stéréochimie R; S; ou RS),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$ , ou  $\text{CH}_2\text{OR}^5$ ; ou  $\text{R}^4$  représente un groupe alkyle, arylalkyle ou aryle en  $\text{C}_4$  à  $\text{C}_{20}$  qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène à condition que  $\text{R}^1$  et  $\text{R}^2$  ne représentent pas simultanément le groupe alkoxy;

$\text{R}^5$  représente un groupe alkyle, aryle ou arylalkyle en  $\text{C}_1$  à  $\text{C}_{20}$  qui peut être substitué ou non substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci; à condition que soient exclus les composés de la formule I ci-dessus dans lesquels :


B représente l'adénine, la xanthine, l'hypoxanthine, la guanine, la 2-aminopurine, la 2,6-diaminopurine, la cytosine, la thymine, ou l'uracile,



$R^3$  représente

où  $alk_1$  est lié à B,  $alk_1$ ,  $alk_2$  et  $alk_3$  sont indépendamment choisis parmi une liaison chimique ou un groupe alkylène en  $C_1$  à  $C_4$ ,  $R^a$  représente l'hydrogène ou un groupe alkyle en  $C_1$  à  $C_4$  et  $Q^1$  représente l'hydrogène ou un groupe hydroxyle, et  $R^1$  et  $R^2$  représentent des groupes alkoxy en  $C_4$  à  $C_6$ , phénoxy ou phényl- ( $C_1$  à  $C_4$ )-alkoxy non substitués.

2. Composé selon la revendication 1 qui a la formule développée générale II :




dans laquelle

B,  $R^1$  et  $R^2$  sont tels que décrits dans la revendication 1, à condition que lorsque Q représente  $NCHN(R^5)_2$ , alors  $R^5$  ne représente pas  $CH_3$ ;

X représente l'hydrogène,  $CH_2OR^6$  (stéréochimie R; S; ou RS), ou un groupe alkyle inférieur substitué ou non substitué, en particulier le groupe méthyle ou hydroxyméthyle; lorsque X représente  $CH_2OR^6$ ,  $R^1$  et  $R^2$  peuvent en outre être indépendamment choisis parmi OH; et

$R^6$  représente un groupe ester hydrolysable.

3. Composé ayant la formule développée générale III :



dans laquelle

B, et  $R^1$  sont tels que précédemment décrits dans la revendication 1;

X représente l'hydrogène;  $CH_2OR^6$  (stéréochimie R; S; ou RS) ou un groupe alkyle inférieur substitué ou non

substitué, en particulier le groupe méthyle ou hydroxyméthyle; lorsque X représente  $\text{CH}_2\text{OR}^6$ ,  $\text{R}^1$  peut en outre représenter OH; et  $\text{R}^6$  représente un groupe ester hydrolysable;

$\text{R}^7$  représente OH,  $\text{NH}_2$ ,  $\text{NHR}^5$ , ou  $\text{NR}^5_2$ ; et

$\text{R}^5$  est tel que décrit dans la revendication 1; sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci;


à condition que soient exclus les composés de la formule III ci-dessus dans laquelle :

B représente l'adénine, la xanthine, l'hypoxanthine, la guanine, la 2-aminopurine, la 2,6-diaminopurine, la cytosine, la thymine, ou l'uracile,

X représente  $\text{alk}_2\text{-Q}^1$

où  $\text{alk}_2$  est choisi parmi une liaison chimique ou un groupe alkylène en  $\text{C}_1$  à  $\text{C}_4$ ,  $\text{Q}^1$  représente l'hydrogène ou le groupe hydroxyle, et  $\text{R}^1$  et  $\text{R}^7$  représentent OH, un groupe alkoxy en  $\text{C}_4$  à  $\text{C}_6$ , phénoxy ou phényl- ( $\text{C}_1$  à  $\text{C}_4$ )-alkoxy non substitué.

4. Composé selon la revendication 1 qui a la formule développée générale IV :



dans laquelle

$\text{R}^8$  et  $\text{R}^9$  sont identiques ou différents et représentent indépendamment  $\text{NR}^{12}$ , ou l'oxygène;

$\text{R}^{10}$  et  $\text{R}^{11}$  sont identiques ou différents et représentent indépendamment l'hydrogène, ou  $\text{R}^5$ ;


$\text{R}^{12}$  représente l'hydrogène ou un groupe alkyle inférieur;

m et n sont identiques ou différents et représentent indépendamment 0 ou 1;

B et  $\text{R}^5$  sont tels que décrits dans la revendication 1; et

X est tel que décrit dans la revendication 2.

5. Composé qui a la formule développée générale V :



\* stéréochimie R; S; ou RS

dans laquelle

$\text{R}^{13}$  représente  $\text{OR}^4$ ,  $\text{NHR}^5$ ,  $\text{NR}^5_2$ , ou OH, à condition que  $\text{R}^{13}$  ne représente pas OH lorsque B représente A

ou C; et

B, R<sup>4</sup>, et R<sup>5</sup> sont tels que décrits dans la revendication 1;  
sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.

5

6. Composé selon la revendication 1, dans lequel R<sup>4</sup> représente un groupe alkyle, aryle ou aryl-alkyle en C<sub>4</sub> à C<sub>20</sub> qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi les groupes hydroxy et halogène.

10

7. Composé selon la revendication 1 dans lequel R<sup>1</sup> et R<sup>2</sup> représentent OR<sup>4</sup>, et R<sup>4</sup> représente un groupe ester physiologiquement hydrolysable choisi parmi CH<sub>2</sub>OC(O)R<sup>5</sup> et CH(R<sup>5</sup>)OC(O)R<sup>5</sup> (stéréochimie R; S; ou RS); ou le groupe aryle dans lequel le groupe aryle est substitué par des substituants indépendamment choisis parmi les groupes hydroxy et halogène.

15

8. Composé selon la revendication 1 qui est

di-(ester propionyloxyméthylque) de PMEA;

di-(ester isobutyryloxyméthylque) de PMEA;

20

bis-diéthylamide de PMEA;

di-(ester acétate de butyle) de PMEA;

25

di-(ester acétate d'éthyle) de PMEA;

di-(ester benzyloxyméthylque) de PMEA;

dibutylamide de PMEA;

30

di-(ester 2-méthylpropylque) de PMEA; ou

di-(ester 3-méthylbutylque) de PMEA.

35

9. Composé selon la revendication 1 qui est

di-(ester pivaloyloxyméthylque) de PMEA.

40

10. Composé selon la revendication 3 qui est

mono-(ester pivaloyloxyméthylque) de PMEA;

mono-(ester octylque) de PMEA;

45

mono-(ester hydroxy-2,2-diméthylpropylque) de PMEA;

mono-(ester 2,2,2-trichloroéthylque) de PMEA; ou

mono-(ester 2,2-difluoro-3-hydroxypropylque) de PMEA.

50

11. Composé selon la revendication 4 qui est

diester cyclopropanylique de PMEA;

55

diester (2,2-diméthyl)cyclopropanylique de PMEA;

diamide (2,2-diméthyl)cyclopropanylique de PMEA

diamide N,N'-diméthyl-cyclopropanylique de PMEÀ.

12. Composé qui est

di-(ester isopropylique) de PMEÀ;

mono-(ester isopropylique) de PMEÀ;

di-(ester isopropylique) de 6-chloro-9-(2-phosphonylméthoxy)éthylpurine;

di-(ester isopropylique) de 9-(2-phosphonylméthoxy)éthylpurine;

mono-(ester isopropylique) de PMEÀ;

ester (mono-isopropylique, mono-pivaloyloxyméthylque) de PMEÀ; ou

ester (mono-isopropylique, mono-phénylique) de PMEÀ;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.

13. Composé qui est

di-(ester phénylique) de PMEÀ;

di-(ester p-nitrobenzylique) de PMEÀ;

di-(ester 2,2,2-trichloroéthylque) de PMEÀ;

di-(ester p-trifluorométhylbenzylique) de PMEÀ;

mono-(ester cholinique) de PMEÀ;

ester (mono-N,N-diéthylacétamidique, mono-pyvaloyloxyméthylque) de PMEÀ;

mono-(ester 3-hydroxypropanylique) de PMEÀ;

mono-(3-amino-2,2-diméthylpropylamide) de PMEÀ;

mono-(ester N,N-diéthylacétamidique) de PMEÀ;

mono-(ester de l'acide acétique) de PMEÀ;

mono-(ester N,N-diisopropylacétamidique) de PMEÀ;

mono-(ester p-nitrobenzylique) de PMEÀ; ou

mono-(ester p-trifluorométhylbenzylique) de PMEÀ;

mono-(ester phénylique) de PMEÀ;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.

14. Procédé de production du composé selon la revendication 1, 2, 12 ou 13 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'alkylation du phosphonate avec l'halogénure d'alkyle approprié; ou l'hydrolyse des diéthers ou des diamines.


15. Procédé de production du composé selon la revendication 3, 12 ou 13 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'hydrolyse des diéthers ou des diamines.



16. Procédé de production du composé selon la revendication 4 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié.
17. Procédé de production du composé selon la revendication 5 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'alkylation du phosphonate avec l'halogénure d'alkyle approprié.
18. Utilisation d'un composé tel que défini dans l'une quelconque des revendications 1 à 13 pour préparer une composition pharmaceutique pour le traitement d'une infection virale chez un mammifère.
19. Utilisation d'un composé tel que défini dans l'une quelconque des revendications 1 à 13 pour préparer une composition pharmaceutique pour inhiber la croissance d'une tumeur chez un mammifère.
20. Composition pharmaceutique qui comprend au moins un composé selon les revendications 1 à 13 associé avec un vecteur ou un excipient sensiblement non toxique pharmaceutiquement acceptable.

#### Revendications pour l'Etat contractant suivant : ES


1. Procédé de production d'un composé ayant la formule développée I :



FORMULE I

dans laquelle

B représente l'adénine (A), la cytosine (C), la guanine (G); la thymine (T), l'uracile (U), la 2,6-diaminopurine (DAP), l'hypoxanthine (Hx),



dans laquelle

Q est indépendamment choisi parmi H, Cl,  $\text{NHR}^5$ ,  $\text{NHR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH ou  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  et  $\text{R}^2$  sont identiques ou différents et représentent indépendamment  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  ou  $\text{N(R}^5)_2$ ;  $\text{R}^1$  et  $\text{R}^2$  étant éventuellement liés l'un à l'autre pour former un groupe cyclique, ou  $\text{R}^1$  ou  $\text{R}^2$  étant éventuellement liés à  $\text{R}^3$  pour former un groupe cyclique;

$\text{R}^3$  représente un groupe alkylène en  $\text{C}_1$  à  $\text{C}_{20}$  qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène; ou  $\text{R}^3$  représente  $\text{CH}(\text{CH}_2\text{OR}^6)\text{CH}_2$ , auquel cas  $\text{R}^1$  et  $\text{R}^2$  peuvent en outre représenter indépendamment OH, et  $\text{R}^6$  représente un groupe ester hydrolysable;

$\text{R}^4$  représente un groupe ester physiologiquement hydrolysable choisi parmi  $\text{CH}_2\text{C(O)NR}^5_2$ ,  $\text{CH}_2\text{C(O)OR}^5$ ,  $\text{CH}_2\text{OC(O)R}^5$ ,  $\text{CH(R}^5\text{)OC(O)R}^5$  (stéréochimie R; S; ou RS),  $\text{CH}_2\text{C(R}^5)_2\text{CH}_2\text{OH}$ , ou  $\text{CH}_2\text{OR}^5$ ; ou  $\text{R}^4$  repré-


sente un groupe alkyle, arylalkyle ou aryle en C<sub>4</sub> à C<sub>20</sub> qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène à condition que R<sup>1</sup> et R<sup>2</sup> ne représentent pas simultanément le groupe alkoxy;

R<sup>5</sup> représente un groupe alkyle, aryle ou arylalkyle en C<sub>1</sub> à C<sub>20</sub> qui peut être substitué ou non substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci; à condition que soient exclus les composés de la formule I ci-dessus dans lesquels :

B représente l'adénine, la xanthine, l'hypoxanthine, la guanine, la 2-aminopurine, la 2,6-diaminopurine, la cytosine, la thymine, ou l'uracile,


R<sup>3</sup> représente



où alk<sub>1</sub> est lié à B, alk<sub>1</sub>, alk<sub>2</sub> et alk<sub>3</sub> sont indépendamment choisis parmi une liaison chimique ou un groupe alkylène en C<sub>1</sub> à C<sub>4</sub>, R<sup>a</sup> représente l'hydrogène ou un groupe alkyle en C<sub>1</sub> à C<sub>4</sub> et Q<sup>1</sup> représente l'hydrogène ou un groupe hydroxyle, et R<sup>1</sup> et R<sup>2</sup> représentent des groupes alkoxy en C<sub>4</sub> à C<sub>6</sub>, phénoxy ou phényl-(C<sub>1</sub> à C<sub>4</sub>)-alkoxy non substitués.

lequel procédé comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'alkylation du phosphonate avec l'halogénure d'alkyle approprié; ou l'hydrolyse des diéthers ou des diamines.

2. Procédé selon la revendication 1 pour produire un composé selon la revendication 1 qui a la formule développée générale II :



FORMULE II


dans laquelle

B, R<sup>1</sup> et R<sup>2</sup> sont tels que décrits dans la revendication 1, à condition que lorsque Q représente NCHN(R<sup>5</sup>)<sub>2</sub>, alors R<sup>5</sup> ne représente pas CH<sub>3</sub>;

X représente l'hydrogène, CH<sub>2</sub>OR<sup>6</sup> (stéréochimie R; S; ou RS), ou un groupe alkyle inférieur substitué ou non substitué, en particulier le groupe méthyle ou hydroxyméthyle; lorsque X représente CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> et R<sup>2</sup> peuvent en outre être indépendamment choisis parmi OH; et

R<sup>6</sup> représente un groupe ester hydrolysable.

3. Procédé de production d'un composé ayant la formule développée générale III :



FORMULE III

dans laquelle

B, et R<sup>1</sup> sont tels que précédemment décrits dans la revendication 1;

X représente l'hydrogène, CH<sub>2</sub>OR<sup>6</sup> (stéréochimie R; S; ou RS) ou un groupe alkyle inférieur substitué ou non substitué, en particulier le groupe méthyle ou hydroxyméthyle; lorsque X représente CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> peut en outre représenter OH; et R<sup>6</sup> représente un groupe ester hydrolysable;

R<sup>7</sup> représente OH, NH<sub>2</sub>, NHR<sup>5</sup>, ou NR<sup>5</sup><sub>2</sub>; et

R<sup>5</sup> est tel que décrit dans la revendication 1;  
sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci;  
à condition que soient exclus les composés de la formule III ci-dessus dans laquelle :


B représente l'adénine, la xanthine, l'hypoxanthine, la guanine, la 2-aminopurine, la 2,6-diaminopurine, la cytosine, la thymine, ou l'uracile,

X représente alk<sub>2</sub>-Q<sup>1</sup>

où alk<sub>2</sub> est choisi parmi une liaison chimique ou un groupe alkylène en C<sub>1</sub> à C<sub>4</sub>, Q<sup>1</sup> représente l'hydrogène ou le groupe hydroxyle, et R<sup>1</sup> et R<sup>7</sup> représentent OH, un groupe alkoxy en C<sub>4</sub> à C<sub>6</sub>, phénoxy ou phényl-(C<sub>1</sub> à C<sub>4</sub>)-alkoxy non substitué.

lequel procédé comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'hydrolyse des diéthers ou des diamines.

4. Procédé de production d'un composé selon la revendication 1 qui a la formule développée générale IV :



FORMULE IV

dans laquelle

R<sup>8</sup> et R<sup>9</sup> sont identiques ou différents et représentent indépendamment NR<sup>12</sup>, ou l'oxygène;

R<sup>10</sup> et R<sup>11</sup> sont identiques ou différents et représentent indépendamment l'hydrogène, ou R<sup>5</sup>;

R<sup>12</sup> représente l'hydrogène ou un groupe alkyle inférieur;


m et n sont identiques ou différents et représentent indépendamment 0 ou 1;

B et R<sup>5</sup> sont tels que décrits dans la revendication 1; et

X est tel que décrit dans la revendication 2;

le quel procédé comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié.

5. Procédé de production d'un composé qui a la formule développée générale V :



\* stéréochimie R; S; ou RS

FORMULE V

dans laquelle

R<sup>13</sup> représente OR<sup>4</sup>, NHR<sup>5</sup>, NR<sup>5</sup><sub>2</sub>, ou OH, à condition que R<sup>13</sup> ne représente pas OH lorsque B représente A ou C; et

B, R<sup>4</sup>, et R<sup>5</sup> sont tels que décrits dans la revendication 1;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci;

lequel procédé comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'alkylation du phosphonate avec l'halogénure d'alkyle approprié.

6. Procédé selon la revendication 1 pour produire un composé selon la revendication 1, dans lequel R<sup>4</sup> représente un groupe alkyle, aryle ou aryl-alkyle en C<sub>4</sub> à C<sub>20</sub> qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi les groupes hydroxy et halogène.

7. Procédé selon la revendication 1 pour produire un composé selon la revendication 1 dans lequel R<sup>1</sup> et R<sup>2</sup> représentent OR<sup>4</sup>, et R<sup>4</sup> représente un groupe ester physiologiquement hydrolysable choisi parmi CH<sub>2</sub>OC(O)R<sup>5</sup> et CH(R<sup>5</sup>)OC(O)R<sup>5</sup> (stéréochimie R; S; ou RS); ou le groupe aryle dans lequel le groupe aryle est substitué par des substituants indépendamment choisis parmi les groupes hydroxy et halogène.

8. Procédé selon la revendication 1 pour produire un composé selon la revendication 1 qui est

di-(ester propionyloxyméthylque) de PMEa;

di-(ester isobutyryloxyméthylque) de PMEa;

bis-diéthylamide de PMEa;

di-(ester acétate de butyle) de PMEa;

di-(ester acétate d'éthyle) de PMEa;

di-(ester benzyloxyméthylque) de PMEa;

dibutylamide de PMEa;

di-(ester 2-méthylpropylque) de PMEa; ou

di-(ester 3-méthylbutylque) de PMEa.

9. Procédé selon la revendication 1 pour produire un composé selon la revendication 1 qui est

di-(ester pivaloyloxyméthylque) de PMEa.

10. Procédé selon la revendication 3 pour produire un composé selon la revendication 3 qui est

- mono-(ester pivaloyloxyméthylque) de PMEA;
  - mono-(ester octylique) de PMEA;
  - 5 mono-(ester hydroxy-2,2-diméthylpropylique) de PMEA;
  - mono-(ester 2,2,2-trichloroéthylque) de PMEA; ou
  - 10 mono-(ester 2,2-difluoro-3-hydroxypropylique) de PMEA.
11. Procédé selon la revendication 4 pour produire un composé selon la revendication 4 qui est
- diester cyclopropanylique de PMEA;
  - 15 diester (2,2-diméthyl)cyclopropanylique de PMEA;
  - diamide (2,2-diméthyl)cyclopropanylique de PMEA
  - diamide N,N'-diméthyl-cyclopropanylique de PMEA.
  - 20
12. Procédé selon la revendication 1 ou la revendication 3 pour produire un composé qui est
- di-(ester isopropylique) de PMEA;
  - 25 mono-(ester isopropylique) de PMEHx;
  - di-(ester isopropylique) de 6-chloro-9-(2-phosphonylméthoxy)éthylpurine;
  - di-(ester isopropylique) de 9-(2-phosphonylméthoxy)éthylpurine;
  - 30 mono-(ester isopropylique) de PMEA;
  - ester (mono-isopropylique, mono-pivaloyloxyméthylque) de PMEA; ou
  - 35 ester (mono-isopropylique, mono-phénylique) de PMEA;
  - sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.
13. Procédé selon la revendication 1 ou la revendication 3 pour produire un composé qui est
- 40 di-(ester phénylique) de PMEA;
  - di-(ester p-nitrobenzylique) de PMEA;
  - di-(ester 2,2,2-trichloroéthylque) de PMEA;
  - 45 di-(ester p-trifluorométhylbenzylique) de PMEA;
  - mono-(ester cholinique) de PMEA;
  - 50 ester (mono-N,N-diéthylacétamidique, mono-pivaloyloxyméthylque) de PMEA;
  - mono-(ester 3-hydroxypropanylique) de PMEA;
  - mono-(3-amino-2,2-diméthylpropylamide) de PMEA;
  - 55 mono-(ester N,N-diéthylacétamidique) de PMEA;
  - mono-(ester de l'acide acétique) de PMEA;

mono-(ester N,N-diisopropylacétamidique) de PME A;

mono-(ester p-nitrobenzylique) de PME A; ou

mono-(ester p-trifluorométhylbenzylique) de PME A;

mono-(ester phénylique) de PME A;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.


14. Utilisation d'un composé tel que défini dans l'une quelconque des revendications 1 à 13 pour préparer une composition pharmaceutique pour le traitement d'une infection virale chez un mammifère.

15. Utilisation d'un composé tel que défini dans l'une quelconque des revendications 1 à 13 pour préparer une composition pharmaceutique pour inhiber la croissance d'une tumeur chez un mammifère.

16. Procédé de préparation d'une composition pharmaceutique qui comprend le mélange d'une proportion d'au moins un composé tel que défini dans l'une quelconque des revendications 1 à 13 avec un vecteur ou un excipient sensiblement non toxique pharmaceutiquement acceptable.

Revendications pour l'Etat contractant suivant : GR


1. Composé ayant la formule développée I :



FORMULE I

dans laquelle

B représente l'adénine (A), la cytosine (C), la guanine (G), la thymine (T), l'uracile (U), la 2,6-diaminopurine (DAP), l'hypoxanthine (Hx),



dans laquelle

Q est indépendamment choisi parmi H, Cl,  $\text{NHR}^5$ ,  $\text{NHR}^5_2$ ,  $\text{NHC(O)R}^5$ ,  $\text{N(C(O)R}^5)_2$ , OH ou  $\text{NCHN(R}^5)_2$ ;

$\text{R}^1$  et  $\text{R}^2$  sont identiques ou différents et représentent indépendamment  $\text{OR}^4$ ,  $\text{NH}_2$ ,  $\text{NHR}^5$  ou  $\text{N(R}^5)_2$ ;  $\text{R}^1$  et  $\text{R}^2$  étant éventuellement liés l'un à l'autre pour former un groupe cyclique, ou  $\text{R}^1$  ou  $\text{R}^2$  étant éventuellement liés à  $\text{R}^3$  pour former un groupe cyclique;


$\text{R}^3$  représente un groupe alkylène en  $\text{C}_1$  à  $\text{C}_{20}$  qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène; ou  $\text{R}^3$  représente  $\text{CH(CH}_2\text{OR}^5)_2$ , auquel cas  $\text{R}^1$  et  $\text{R}^2$  peuvent en outre représenter indépendamment OH, et  $\text{R}^5$  représente un groupe ester hydrolysable;

$R^4$  représente un groupe ester physiologiquement hydrolysable choisi parmi  $\text{CH}_2\text{C}(\text{O})\text{NR}^5$ ,  $\text{CH}_2\text{C}(\text{O})\text{OR}^5$ ,  $\text{CH}_2\text{OC}(\text{O})\text{R}^5$ ,  $\text{CH}(\text{R}^5)\text{OC}(\text{O})\text{R}^5$  (stéréochimie R; S; ou RS),  $\text{CH}_2\text{C}(\text{R}^5)_2\text{CH}_2\text{OH}$ , ou  $\text{CH}_2\text{OR}^5$ ; ou  $R^4$  représente un groupe alkyle, arylalkyle ou aryle en  $\text{C}_4$  à  $\text{C}_{20}$  qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène à condition que  $R^1$  et  $R^2$  ne représentent pas simultanément le groupe alkoxy;

$R^5$  représente un groupe alkyle, aryle ou arylalkyle en  $\text{C}_1$  à  $\text{C}_{20}$  qui peut être substitué ou non substitué par des substituants indépendamment choisis parmi le groupe constitué du groupe hydroxy et du groupe halogène;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci; à condition que soient exclus les composés de la formule I ci-dessus dans lesquels :


B représente l'adénine, la xanthine, l'hypoxanthine, la guanine, la 2-aminopurine, la 2,6-diaminopurine, la cytosine, la thymine, ou l'uracile,



$R^3$  représente

où  $\text{alk}_1$  est lié à B,  $\text{alk}_1$ ,  $\text{alk}_2$  et  $\text{alk}_3$  sont indépendamment choisis parmi une liaison chimique ou un groupe alkylène en  $\text{C}_1$  à  $\text{C}_4$ ,  $R^a$  représente l'hydrogène ou un groupe alkyle en  $\text{C}_1$  à  $\text{C}_4$  et  $Q^1$  représente l'hydrogène ou un groupe hydroxyle, et  $R^1$  et  $R^2$  représentent des groupes alkoxy en  $\text{C}_4$  à  $\text{C}_6$ , phénoxy ou phényl- ( $\text{C}_1$  à  $\text{C}_4$ )-alkoxy non substitués.

2. Composé selon la revendication 1 qui a la formule développée générale II :



FORMULE II


dans laquelle

B,  $R^1$  et  $R^2$  sont tels que décrits dans la revendication 1, à condition que lorsque Q représente  $\text{NCHN}(\text{R}^5)_2$ , alors  $R^5$  ne représente pas  $\text{CH}_3$ ;

X représente l'hydrogène,  $\text{CH}_2\text{OR}^6$  (stéréochimie R; S; ou RS), ou un groupe alkyle inférieur substitué ou non substitué, en particulier le groupe méthyle ou hydroxyméthyle; lorsque X représente  $\text{CH}_2\text{OR}^6$ ,  $R^1$  et  $R^2$  peuvent en outre être indépendamment choisis parmi OH; et

$R^6$  représente un groupe ester hydrolysable.

3. Composé ayant la formule développée générale III :



FORMULE III

dans laquelle

B, et R<sup>1</sup> sont tels que précédemment décrits dans la revendication 1;

X représente l'hydrogène, CH<sub>2</sub>OR<sup>6</sup> (stéréochimie R; S; ou RS) ou un groupe alkyle inférieur substitué ou non substitué, en particulier le groupe méthyle ou hydroxyméthyle; lorsque X représente CH<sub>2</sub>OR<sup>6</sup>, R<sup>1</sup> peut en outre représenter OH; et R<sup>6</sup> représente un groupe ester hydrolysable;

R<sup>7</sup> représente OH, NH<sub>2</sub>, NHR<sup>5</sup>, ou NR<sup>5</sup><sub>2</sub>; et


R<sup>5</sup> est tel que décrit dans la revendication 1;  
sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci;  
à condition que soient exclus les composés de la formule III ci-dessus dans laquelle :

B représente l'adénine, la xanthine, l'hypoxanthine, la guanine, la 2-aminopurine, la 2,6-diaminopurine, la cytosine, la thymine, ou l'uracile,

X représente alk<sub>2</sub>-Q<sup>1</sup>

où alk<sub>2</sub> est choisi parmi une liaison chimique ou un groupe alkylène en C<sub>1</sub> à C<sub>4</sub>, Q<sup>1</sup> représente l'hydrogène ou le groupe hydroxyle, et R<sup>1</sup> et R<sup>7</sup> représentent OH, un groupe alkoxy en C<sub>4</sub> à C<sub>6</sub>, phénoxy ou phényl-(C<sub>1</sub> à C<sub>4</sub>)-alkoxy non substitué.

4. Composé selon la revendication 1 qui a la formule développée générale IV :



FORMULE IV

dans laquelle

R<sup>8</sup> et R<sup>9</sup> sont identiques ou différents et représentent indépendamment NR<sup>12</sup>, ou l'oxygène;

R<sup>10</sup> et R<sup>11</sup> sont identiques ou différents et représentent indépendamment l'hydrogène, ou R<sup>5</sup>;

R<sup>12</sup> représente l'hydrogène ou un groupe alkyle inférieur;


m et n sont identiques ou différents et représentent indépendamment 0 ou 1;

B et R<sup>5</sup> sont tels que décrits dans la revendication 1; et

X est tel que décrit dans la revendication 2.



5. Composé qui a la formule développée générale V :



\* stéréochimie R; S; ou RS

FORMULE V

dans laquelle

- 15  $R^{13}$  représente  $OR^4$ ,  $NHR^5$ ,  $NR^5_2$ , ou OH, à condition que  $R^{13}$  ne représente pas OH lorsque B représente A ou C; et

B,  $R^4$ , et  $R^5$  sont tels que décrits dans la revendication 1;  
sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.

- 20 6. Composé selon la revendication 1, dans lequel  $R^4$  représente un groupe alkyle, aryle ou aryl-alkyle en  $C_4$  à  $C_{20}$  qui peut être non substitué ou substitué par des substituants indépendamment choisis parmi les groupes hydroxy et halogène.
- 25 7. Composé selon la revendication 1 dans lequel  $R^1$  et  $R^2$  représentent  $OR^4$ , et  $R^4$  représente un groupe ester physiologiquement hydrolysable choisi parmi  $CH_2OC(O)R^5$  et  $CH(R^5)OC(O)R^5$  (stéréochimie R; S; ou RS); ou le groupe aryle dans lequel le groupe aryle est substitué par des substituants indépendamment choisis parmi les groupes hydroxy et halogène.
- 30 8. Composé selon la revendication 1 qui est
- di-(ester propionyloxyméthylrique) de PMEa;
  - di-(ester isobutyryloxyméthylrique) de PMEa;

35 bis-diéthylamide de PMEa;

  - di-(ester acétate de butyle) de PMEa;

40 di-(ester acétate d'éthyle) de PMEa;

  - di-(ester benzoxyloxyméthylrique) de PMEa;
  - dibutylamide de PMEa;

45 di-(ester 2-méthylpropylrique) de PMEa; ou

  - di-(ester 3-méthylbutylrique) de PMEa.
- 50 9. Composé selon la revendication 1 qui est
- di-(ester pivaloxyloxyméthylrique) de PMEa.
- 55 10. Composé selon la revendication 3 qui est
- mono-(ester pivaloxyloxyméthylrique) de PMEa;
  - mono-(ester octylrique) de PMEa;

- mono-(ester hydroxy-2,2-diméthylpropylique) de PMEa;
- mono-(ester 2,2,2-trichloroéthylque) de PMEa; ou
- 5 mono-(ester 2,2-difluoro-3-hydroxypropylique) de PMEa.

11. Composé selon la revendication 4 qui est

- diester cyclopropylique de PMEa;
- 10 diester (2,2-diméthyl)cyclopropylique de PMEa;
- diamide (2,2-diméthyl)cyclopropylique de PMEa
- 15 diamide N,N'-diméthyl-cyclopropylique de PMEa.

12. Composé qui est

- di-(ester isopropylique) de PMEa;
- 20 mono-(ester isopropylique) de PMEhx;
- di-(ester isopropylique) de 6-chloro-9-(2-phosphonilméthoxy)éthylpurine;
- 25 di-(ester isopropylique) de 9-(2-phosphonilméthoxy)éthylpurine;
- mono-(ester isopropylique) de PMEa;
- ester (mono-isopropylique, mono-pivaloyloxyméthylque) de PMEa; ou
- 30 ester (mono-isopropylique, mono-phénylique) de PMEa;
- sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.

13. Composé qui est

- 35 di-(ester phénylique) de PMEa;
- di-(ester p-nitrobenzylique) de PMEa;
- 40 di-(ester 2,2,2-trichloroéthylque) de PMEa;
- di-(ester p-trifluorométhylbenzylique) de PMEa;
- mono-(ester cholinique) de PMEa;
- 45 ester (mono-N,N-diéthylacétamidique, mono-pivaloyloxyméthylque) de PMEa;
- mono-(ester 3-hydroxypropylique) de PMEa;
- 50 mono-(3-amino-2,2-diméthylpropylamide)) de PMEa;
- mono-(ester N,N-diéthylacétamidique) de PMEa;
- mono-(ester de l'acide acétique) de PMEa;
- 55 mono-(ester N,N-diisopropylacétamidique) de PMEa;
- mono-(ester p-nitrobenzylique) de PMEa; ou

mono-(ester p-trifluorométhylbenzylique) de PMEAs;

mono-(ester phénylique) de PMEAs;

sels d'addition d'acide, sels de métal, et solvates pharmaceutiquement acceptables de celui-ci.

5

14. Procédé de production du composé selon la revendication 1, 2, 12 ou 13 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'alkylation du phosphonate avec l'halogénure d'alkyle approprié; ou l'hydrolyse des diéthers ou des diamines.

10

15. Procédé de production du composé selon la revendication 3, 12 ou 13 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'hydrolyse des diéthers ou des diamines.

15

16. Procédé de production du composé selon la revendication 4 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié.

17. Procédé de production du composé selon la revendication 5 qui comprend la réaction du phosphonate avec un agent activant, puis la réaction avec l'amine ou l'alcool approprié; ou l'alkylation du phosphonate avec l'halogénure d'alkyle approprié.

20

18. Utilisation d'un composé tel que défini dans l'une quelconque des revendications 1 à 13 pour préparer une composition pharmaceutique pour le traitement d'une infection virale chez un mammifère.

25

19. Utilisation d'un composé tel que défini dans l'une quelconque des revendications 1 à 13 pour préparer une composition pharmaceutique pour inhiber la croissance d'une tumeur chez un mammifère.

20. Procédé de préparation d'une composition pharmaceutique qui comprend le mélange d'une proportion d'au moins un composé tel que défini dans l'une quelconque des revendications 1 à 13 avec un vecteur ou un excipient sensiblement non toxique pharmaceutiquement acceptable.

30

35

40

45

50

55